

Cultivating Complex Analysis:
Definition (of analytic functions) (2.4.1)
Analytic functions are holomorphic (2.4.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Analytic functions are functions equal to a convergent power series near every point.

Definition

Let $U \subset \mathbb{C}$ be open. A function $f: U \rightarrow \mathbb{C}$ is *analytic* if for every $p \in U$, there exists an $r > 0$ and a power series $\sum c_n(z - p)^n$ converging to f on $\Delta_r(p) \subset U$.

Analytic functions are functions equal to a convergent power series near every point.

Definition

Let $U \subset \mathbb{C}$ be open. A function $f: U \rightarrow \mathbb{C}$ is *analytic* if for every $p \in U$, there exists an $r > 0$ and a power series $\sum c_n(z - p)^n$ converging to f on $\Delta_r(p) \subset U$.

Eventually, we will see that function is holomorphic if and only if it is analytic.

Analytic functions are functions equal to a convergent power series near every point.

Definition

Let $U \subset \mathbb{C}$ be open. A function $f: U \rightarrow \mathbb{C}$ is *analytic* if for every $p \in U$, there exists an $r > 0$ and a power series $\sum c_n(z - p)^n$ converging to f on $\Delta_r(p) \subset U$.

Eventually, we will see that function is holomorphic if and only if it is analytic.

But that is not so easy. Today we will prove that analytic \Rightarrow holomorphic.

Analytic functions are functions equal to a convergent power series near every point.

Definition

Let $U \subset \mathbb{C}$ be open. A function $f: U \rightarrow \mathbb{C}$ is *analytic* if for every $p \in U$, there exists an $r > 0$ and a power series $\sum c_n(z - p)^n$ converging to f on $\Delta_r(p) \subset U$.

Eventually, we will see that function is holomorphic if and only if it is analytic.

But that is not so easy. Today we will prove that analytic \Rightarrow holomorphic.

Remark: A subtle point is that it is not immediate that a convergent power series is analytic.

Proposition

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by

$$f(z) = \sum_{n=0}^{\infty} c_n (z - p)^n, \quad \text{converging in } \Delta_R(p).$$

Proposition

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by

$$f(z) = \sum_{n=0}^{\infty} c_n(z-p)^n, \quad \text{converging in } \Delta_R(p).$$

Then f is complex differentiable at every $z \in \Delta_R(p)$, and

$$f'(z) = \sum_{n=1}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging in } \Delta_R(p).$$

Proposition

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by

$$f(z) = \sum_{n=0}^{\infty} c_n(z-p)^n, \quad \text{converging in } \Delta_R(p).$$

Then f is complex differentiable at every $z \in \Delta_R(p)$, and

$$f'(z) = \sum_{n=1}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging in } \Delta_R(p).$$

Proof: WLOG $p = 0$.

Proposition

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by

$$f(z) = \sum_{n=0}^{\infty} c_n(z-p)^n, \quad \text{converging in } \Delta_R(p).$$

Then f is complex differentiable at every $z \in \Delta_R(p)$, and

$$f'(z) = \sum_{n=1}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging in } \Delta_R(p).$$

Proof: WLOG $p = 0$.

Consider the difference quotient of z^n at z_0

$$\frac{z^n - z_0^n}{z - z_0} = \sum_{k=0}^{n-1} z^k z_0^{n-1-k}$$

Proposition

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by

$$f(z) = \sum_{n=0}^{\infty} c_n(z-p)^n, \quad \text{converging in } \Delta_R(p).$$

Then f is complex differentiable at every $z \in \Delta_R(p)$, and

$$f'(z) = \sum_{n=1}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging in } \Delta_R(p).$$

Proof: WLOG $p = 0$.

Consider the difference quotient of z^n at z_0

$$\frac{z^n - z_0^n}{z - z_0} = \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \xrightarrow{\text{as } z \rightarrow z_0} n z_0^{n-1}.$$

Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0}$$

Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0}$$

Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .

Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 . It is continuous if the series converges uniformly for z in a neighborhood of z_0 .


Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .

It is continuous if the series converges uniformly for z in a neighborhood of z_0 .

Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.



Suppose $z_0, z \in \Delta_R(0)$,


$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .

It is continuous if the series converges uniformly for z in a neighborhood of z_0 .

Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.

$$\left| c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \right| \leq \sum_{k=0}^{n-1} |c_n| s^{n-1} = n |c_n| s^{n-1} = n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}$$



Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$


We need to show that the expression on the right is a continuous function of z at z_0 .

It is continuous if the series converges uniformly for z in a neighborhood of z_0 .

Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.

$$\left| c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \right| \leq \sum_{k=0}^{n-1} |c_n| s^{n-1} = n |c_n| s^{n-1} = n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}$$

$|c_n| r^{n-1}$ is bounded by some $M > 0$ (f converges in $\Delta_R(0)$ and $r < R$)



Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .


It is continuous if the series converges uniformly for z in a neighborhood of z_0 .

Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.

$$\left| c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \right| \leq \sum_{k=0}^{n-1} |c_n| s^{n-1} = n |c_n| s^{n-1} = n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}$$

$|c_n| r^{n-1}$ is bounded by some $M > 0$ (f converges in $\Delta_R(0)$ and $r < R$)

$$\sqrt[n]{n |c_n| s^{n-1}} = \sqrt[n]{n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}} \leq \sqrt[n]{n M \left(\frac{s}{r} \right)^{n-1}} \xrightarrow{\text{as } n \rightarrow \infty} \frac{s}{r} < 1$$



Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .

It is continuous if the series converges uniformly for z in a neighborhood of z_0 .


Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.

$$\left| c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \right| \leq \sum_{k=0}^{n-1} |c_n| s^{n-1} = n |c_n| s^{n-1} = n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}$$

$|c_n| r^{n-1}$ is bounded by some $M > 0$ (f converges in $\Delta_R(0)$ and $r < R$)

$$\sqrt[n]{n |c_n| s^{n-1}} = \sqrt[n]{n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}} \leq \sqrt[n]{n M \left(\frac{s}{r} \right)^{n-1}} \xrightarrow{\text{as } n \rightarrow \infty} \frac{s}{r} < 1$$

so $\sum n |c_n| s^{n-1}$ converges (root test).



Suppose $z_0, z \in \Delta_R(0)$,

$$\frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \frac{z^n - z_0^n}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}.$$

We need to show that the expression on the right is a continuous function of z at z_0 .

It is continuous if the series converges uniformly for z in a neighborhood of z_0 .

Suppose $0 < s < r < R$ and $z_0, z \in \Delta_s(0)$.


$$\left| c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k} \right| \leq \sum_{k=0}^{n-1} |c_n| s^{n-1} = n |c_n| s^{n-1} = n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}$$

$|c_n| r^{n-1}$ is bounded by some $M > 0$ (f converges in $\Delta_R(0)$ and $r < R$)

$$\sqrt[n]{n |c_n| s^{n-1}} = \sqrt[n]{n |c_n| r^{n-1} \left(\frac{s}{r} \right)^{n-1}} \leq \sqrt[n]{n M \left(\frac{s}{r} \right)^{n-1}} \xrightarrow{\text{as } n \rightarrow \infty} \frac{s}{r} < 1$$

so $\sum n |c_n| s^{n-1}$ converges (root test).

So $\sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z^k z_0^{n-1-k}$, converges uniformly in z on $\Delta_s(p)$.



So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z_0^k z_0^{n-1-k} = \sum_{n=1}^{\infty} n c_n z_0^{n-1}.$$

So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z_0^k z_0^{n-1-k} = \sum_{n=1}^{\infty} n c_n z_0^{n-1}.$$

As s and r were arbitrary we get the result at any $z_0 \in \Delta_R(0)$.



So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z_0^k z_0^{n-1-k} = \sum_{n=1}^{\infty} n c_n z_0^{n-1}.$$

As s and r were arbitrary we get the result at any $z_0 \in \Delta_R(0)$. □

Corollary

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by $f(z) = \sum_{n=0}^{\infty} c_n (z - p)^n$, converging in $\Delta_R(p)$.

So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z_0^k z_0^{n-1-k} = \sum_{n=1}^{\infty} n c_n z_0^{n-1}.$$

As s and r were arbitrary we get the result at any $z_0 \in \Delta_R(0)$. □

Corollary

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by $f(z) = \sum_{n=0}^{\infty} c_n (z - p)^n$, converging in $\Delta_R(p)$.

Then f is infinitely complex differentiable in $\Delta_R(p)$, and the k^{th} derivative is given by

$$f^{(k)}(z) = \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) c_n (z-p)^{n-k}, \quad \text{converging in } \Delta_R(p).$$

So we can swap the limits:

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} = \sum_{n=1}^{\infty} c_n \sum_{k=0}^{n-1} z_0^k z_0^{n-1-k} = \sum_{n=1}^{\infty} n c_n z_0^{n-1}.$$

As s and r were arbitrary we get the result at any $z_0 \in \Delta_R(0)$. □

Corollary

Let $f: \Delta_R(p) \rightarrow \mathbb{C}$ be defined by $f(z) = \sum_{n=0}^{\infty} c_n (z - p)^n$, converging in $\Delta_R(p)$.

Then f is infinitely complex differentiable in $\Delta_R(p)$, and the k^{th} derivative is given by

$$f^{(k)}(z) = \sum_{n=k}^{\infty} n(n-1) \cdots (n-k+1) c_n (z - p)^{n-k}, \quad \text{converging in } \Delta_R(p).$$

Furthermore, $c_n = \frac{f^{(n)}(p)}{n!}$.

A consequence is that the power series is unique since the coefficients are unique

$$c_n = \frac{f^{(n)}(p)}{n!}.$$

A consequence is that the power series is unique since the coefficients are unique

$$c_n = \frac{f^{(n)}(p)}{n!}.$$

In fact, c_n depends only on values of f in arbitrarily small neighborhood of p .

A consequence is that the power series is unique since the coefficients are unique

$$c_n = \frac{f^{(n)}(p)}{n!}.$$

In fact, c_n depends only on values of f in arbitrarily small neighborhood of p .

Applied to the analytic functions we get:

Corollary

An analytic function is infinitely complex differentiable, and each derivative is analytic.