

# Cultivating Complex Analysis: Laurent series (4.4 part 2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato



## Theorem (Existence of Laurent series)

*Suppose that  $0 \leq r_1 < r_2 \leq \infty$  and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is holomorphic.*



## Theorem (Existence of Laurent series)

*Suppose that  $0 \leq r_1 < r_2 \leq \infty$  and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is holomorphic. Then there exist unique numbers  $c_n \in \mathbb{C}$  for  $n \in \mathbb{Z}$  such that*

$$f(z) = \sum_{n=-\infty}^{\infty} c_n (z - p)^n,$$

*converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .*



## Theorem (Existence of Laurent series)

Suppose that  $0 \leq r_1 < r_2 \leq \infty$  and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is holomorphic. Then there exist unique numbers  $c_n \in \mathbb{C}$  for  $n \in \mathbb{Z}$  such that

$$f(z) = \sum_{n=-\infty}^{\infty} c_n (z - p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ . The numbers  $c_n$  are given by

$$c_n = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{(z - p)^{n+1}} dz,$$

where  $\gamma$  is any circle of radius  $s$ ,  $r_1 < s < r_2$ , centered at  $p$  oriented counterclockwise.




**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .



**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .






**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :






**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .






**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .

If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .





**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .


Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .

If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .

So  $\Gamma$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ .





**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .


If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .

If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .

So  $\Gamma$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ .

However, when  $z \in \text{ann}(p; s_1, s_2)$ , then  $n(\Gamma; z) = 1$ .





**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .


If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .

So  $\Gamma$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ .

However, when  $z \in \text{ann}(p; s_1, s_2)$ , then  $n(\Gamma; z) = 1$ .

By Cauchy's formula for  $z \in \text{ann}(p; s_1, s_2)$ ,

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$





**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .


If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .

So  $\Gamma$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ .

However, when  $z \in \text{ann}(p; s_1, s_2)$ , then  $n(\Gamma; z) = 1$ .

By Cauchy's formula for  $z \in \text{ann}(p; s_1, s_2)$ ,

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$





**Proof of the theorem:** Choose  $s_1$  and  $s_2$  such that  $r_1 < s_1 < s_2 < r_2$ .

Define the cycle  $\Gamma = \partial\Delta_{s_2}(p) - \partial\Delta_{s_1}(p)$ .

If  $q \in \mathbb{C} \setminus \text{ann}(p; r_1, r_2)$ , then  $n(\Gamma; q) = 0$ :

If  $q$  is in the hole,  $n(\partial\Delta_{s_j}(p); q) = 1$  for  $j = 1, 2$ .

If  $q$  is on the outside,  $n(\partial\Delta_{s_j}(p); q) = 0$  for  $j = 1, 2$ .


So  $\Gamma$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ .

However, when  $z \in \text{ann}(p; s_1, s_2)$ , then  $n(\Gamma; z) = 1$ .

By Cauchy's formula for  $z \in \text{ann}(p; s_1, s_2)$ ,

$$f(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta - \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

We will expand the two integrals separately.





First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .



First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .

We follow the same reasoning as in the proof of the existence of power series.



First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .

We follow the same reasoning as in the proof of the existence of power series.

$$\frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \frac{1}{1 - \frac{z-p}{\zeta-p}} d\zeta$$



First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .

We follow the same reasoning as in the proof of the existence of power series.

$$\begin{aligned}\frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \frac{1}{1 - \frac{z-p}{\zeta-p}} d\zeta \\ &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p}\right)^n d\zeta\end{aligned}$$



First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .

We follow the same reasoning as in the proof of the existence of power series.

$$\begin{aligned}
 \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \frac{1}{1 - \frac{z-p}{\zeta-p}} d\zeta \\
 &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p}\right)^n d\zeta \\
 &= \sum_{n=0}^{\infty} \underbrace{\left( \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta \right)}_{c_n} (z-p)^n.
 \end{aligned}$$



First, if  $\zeta \in \partial\Delta_{s_2}$ , then  $\left|\frac{z-p}{\zeta-p}\right| = \frac{|z-p|}{s_2} < 1$ .

We follow the same reasoning as in the proof of the existence of power series.

$$\begin{aligned}
 \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \frac{1}{1 - \frac{z-p}{\zeta-p}} d\zeta \\
 &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{\zeta - p} \sum_{n=0}^{\infty} \left(\frac{z-p}{\zeta-p}\right)^n d\zeta \\
 &= \sum_{n=0}^{\infty} \underbrace{\left( \frac{1}{2\pi i} \int_{\partial\Delta_{s_2}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta \right)}_{c_n} (z-p)^n.
 \end{aligned}$$

We can swap the series limit with the integral as the convergence is uniform on the circle.



Similarly, if  $\zeta \in \partial\Delta_{s_1}$ , then  $\left|\frac{\zeta-p}{z-p}\right| = \frac{s_1}{|z-p|} < 1$ , so



Similarly, if  $\zeta \in \partial\Delta_{s_1}$ , then  $\left|\frac{\zeta-p}{z-p}\right| = \frac{s_1}{|z-p|} < 1$ , so

$$-\frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \frac{1}{1 - \frac{\zeta-p}{z-p}} d\zeta$$



Similarly, if  $\zeta \in \partial\Delta_{s_1}$ , then  $\left|\frac{\zeta-p}{z-p}\right| = \frac{s_1}{|z-p|} < 1$ , so

$$\begin{aligned} -\frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \frac{1}{1 - \frac{\zeta-p}{z-p}} d\zeta \\ &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \sum_{m=0}^{\infty} \left(\frac{\zeta - p}{z - p}\right)^m d\zeta \end{aligned}$$



Similarly, if  $\zeta \in \partial\Delta_{s_1}$ , then  $\left|\frac{\zeta-p}{z-p}\right| = \frac{s_1}{|z-p|} < 1$ , so

$$\begin{aligned}
 -\frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \frac{1}{1 - \frac{\zeta-p}{z-p}} d\zeta \\
 &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \sum_{m=0}^{\infty} \left(\frac{\zeta - p}{z - p}\right)^m d\zeta \\
 &= \sum_{m=0}^{\infty} \left( \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} f(\zeta)(\zeta - p)^m d\zeta \right) (z - p)^{-m-1}
 \end{aligned}$$



Similarly, if  $\zeta \in \partial\Delta_{s_1}$ , then  $\left|\frac{\zeta-p}{z-p}\right| = \frac{s_1}{|z-p|} < 1$ , so

$$\begin{aligned}
 -\frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{\zeta - z} d\zeta &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \frac{1}{1 - \frac{\zeta-p}{z-p}} d\zeta \\
 &= \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{z - p} \sum_{m=0}^{\infty} \left(\frac{\zeta - p}{z - p}\right)^m d\zeta \\
 &= \sum_{m=0}^{\infty} \left( \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} f(\zeta)(\zeta - p)^m d\zeta \right) (z - p)^{-m-1} \\
 &= \sum_{n=-\infty}^{-1} \underbrace{\left( \frac{1}{2\pi i} \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta \right)}_{c_n} (z - p)^n.
 \end{aligned}$$



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .

Let  $s$  be such that  $r_1 < s < r_2$ .



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .

Let  $s$  be such that  $r_1 < s < r_2$ .

The cycle  $\partial\Delta_s(p) - \partial\Delta_{s_1}(p)$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ , and  $z \mapsto \frac{f(z)}{(z-p)^{n+1}}$  is holomorphic in  $\text{ann}(p; r_1, r_2)$ .



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .

Let  $s$  be such that  $r_1 < s < r_2$ .

The cycle  $\partial\Delta_s(p) - \partial\Delta_{s_1}(p)$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ , and  $z \mapsto \frac{f(z)}{(z-p)^{n+1}}$  is holomorphic in  $\text{ann}(p; r_1, r_2)$ .

Cauchy's theorem says

$$0 = \int_{\partial\Delta_s(p) - \partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta$$



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .

Let  $s$  be such that  $r_1 < s < r_2$ .

The cycle  $\partial\Delta_s(p) - \partial\Delta_{s_1}(p)$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ , and  $z \mapsto \frac{f(z)}{(z-p)^{n+1}}$  is holomorphic in  $\text{ann}(p; r_1, r_2)$ .

Cauchy's theorem says

$$0 = \int_{\partial\Delta_s(p) - \partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta = \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta - \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta.$$



We got the series, although perhaps not the formula for  $c_n$  for circle of any radius  $s$ .

Let  $s$  be such that  $r_1 < s < r_2$ .

The cycle  $\partial\Delta_s(p) - \partial\Delta_{s_1}(p)$  is homologous to zero in  $\text{ann}(p; r_1, r_2)$ , and  $z \mapsto \frac{f(z)}{(z-p)^{n+1}}$  is holomorphic in  $\text{ann}(p; r_1, r_2)$ .

Cauchy's theorem says

$$0 = \int_{\partial\Delta_s(p) - \partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta = \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta - \int_{\partial\Delta_{s_1}(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta.$$

Similarly for  $s_2$  and so

$$c_n = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta - p)^{n+1}} d\zeta.$$



Next, convergence.



Next, convergence.

For any  $\epsilon > 0$ , the geometric series used for the first part converges uniformly absolutely when  $\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{s_2} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\Delta_{s_2}(p)$ .



Next, convergence.

For any  $\epsilon > 0$ , the geometric series used for the first part converges uniformly absolutely when  $\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{s_2} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\Delta_{s_2}(p)$ .

The geometric series used for the second part converges uniformly absolutely when  $\left| \frac{\zeta-p}{z-p} \right| = \frac{s_1}{|z-p|} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\mathbb{C} \setminus \overline{\Delta_{s_1}(p)}$ .



Next, convergence.

For any  $\epsilon > 0$ , the geometric series used for the first part converges uniformly absolutely when  $\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{s_2} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\Delta_{s_2}(p)$ .

The geometric series used for the second part converges uniformly absolutely when  $\left| \frac{\zeta-p}{z-p} \right| = \frac{s_1}{|z-p|} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\mathbb{C} \setminus \overline{\Delta_{s_1}(p)}$ .

Any compact subset of  $\text{ann}(p; r_1, r_2)$  is a compact inside both of these for some  $s_1, s_2$ .



Next, convergence.

For any  $\epsilon > 0$ , the geometric series used for the first part converges uniformly absolutely when  $\left| \frac{z-p}{\zeta-p} \right| = \frac{|z-p|}{s_2} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\Delta_{s_2}(p)$ .

The geometric series used for the second part converges uniformly absolutely when  $\left| \frac{\zeta-p}{z-p} \right| = \frac{s_1}{|z-p|} \leq 1 - \epsilon$ .

So the resulting series converges uniformly absolutely on compact subsets of  $\mathbb{C} \setminus \overline{\Delta_{s_1}(p)}$ .

Any compact subset of  $\text{ann}(p; r_1, r_2)$  is a compact inside both of these for some  $s_1, s_2$ .

So the full series converges uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .



Finally, uniqueness.



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .

Then



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .

Then

$$c_m = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta-p)^{m+1}} d\zeta$$



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .

Then

$$c_m = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta-p)^{m+1}} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \left( \sum_{n=-\infty}^{\infty} d_n(\zeta-p)^n \right) \frac{1}{(\zeta-p)^{m+1}} d\zeta$$



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .

Then

$$\begin{aligned} c_m &= \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta-p)^{m+1}} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \left( \sum_{n=-\infty}^{\infty} d_n(\zeta-p)^n \right) \frac{1}{(\zeta-p)^{m+1}} d\zeta \\ &= \frac{1}{2\pi i} \sum_{n=-\infty}^{\infty} d_n \int_{\partial\Delta_s(p)} (\zeta-p)^{n-m-1} d\zeta \end{aligned}$$



Finally, uniqueness.

Suppose

$$f(z) = \sum_{n=-\infty}^{\infty} d_n(z-p)^n,$$

converging uniformly absolutely on compact subsets of  $\text{ann}(p; r_1, r_2)$ .

Then

$$\begin{aligned} c_m &= \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \frac{f(\zeta)}{(\zeta-p)^{m+1}} d\zeta = \frac{1}{2\pi i} \int_{\partial\Delta_s(p)} \left( \sum_{n=-\infty}^{\infty} d_n(\zeta-p)^n \right) \frac{1}{(\zeta-p)^{m+1}} d\zeta \\ &= \frac{1}{2\pi i} \sum_{n=-\infty}^{\infty} d_n \int_{\partial\Delta_s(p)} (\zeta-p)^{n-m-1} d\zeta = d_m. \end{aligned}$$

(The last equality because  $\int_{\partial\Delta_s(p)} (\zeta-p)^{n-m-1} d\zeta \neq 0$  only when  $n = m$ .)

□



We differentiate/antidifferentiate formally (except for antidifferentiating the  $c_{-1}(z - p)^{-1}$ ).



We differentiate/antidifferentiate formally (except for antidifferentiating the  $c_{-1}(z-p)^{-1}$ ).

### Proposition

Suppose  $p \in \mathbb{C}$ ,  $0 \leq r_1 < r_2 \leq \infty$ , and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is defined by

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$



We differentiate/antidifferentiate formally (except for antidifferentiating the  $c_{-1}(z-p)^{-1}$ ).

### Proposition

Suppose  $p \in \mathbb{C}$ ,  $0 \leq r_1 < r_2 \leq \infty$ , and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is defined by

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

(i) Then  $f$  is holomorphic and its derivative is defined by

$$f'(z) = \sum_{n=-\infty}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$



We differentiate/antidifferentiate formally (except for antidifferentiating the  $c_{-1}(z-p)^{-1}$ ).

## Proposition

Suppose  $p \in \mathbb{C}$ ,  $0 \leq r_1 < r_2 \leq \infty$ , and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is defined by

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

(i) Then  $f$  is holomorphic and its derivative is defined by

$$f'(z) = \sum_{n=-\infty}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

(ii) If  $c_{-1} = 0$ , then  $f = F'$ , where

$$F(z) = \sum_{n=-\infty, n \neq -1}^{\infty} \frac{c_n}{n+1} (z-p)^{n+1}, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$



We differentiate/antidifferentiate formally (except for antidifferentiating the  $c_{-1}(z-p)^{-1}$ ).

### Proposition

Suppose  $p \in \mathbb{C}$ ,  $0 \leq r_1 < r_2 \leq \infty$ , and  $f: \text{ann}(p; r_1, r_2) \rightarrow \mathbb{C}$  is defined by

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z-p)^n, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

(i) Then  $f$  is holomorphic and its derivative is defined by

$$f'(z) = \sum_{n=-\infty}^{\infty} n c_n(z-p)^{n-1}, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

(ii) If  $c_{-1} = 0$ , then  $f = F'$ , where

$$F(z) = \sum_{n=-\infty, n \neq -1}^{\infty} \frac{c_n}{n+1} (z-p)^{n+1}, \quad \text{converging uniformly on compact subsets of } \text{ann}(p; r_1, r_2).$$

**Proof:** Exercise.



Some more useful exercises:



Some more useful exercises:

**Exercise:** Suppose  $f$  and  $g$  are holomorphic functions defined on  $\text{ann}(p; r_1, r_2)$ . Let  $a_n$  be the coefficients in the Laurent series for  $f$  and  $b_n$  be the coefficients in the Laurent series for  $g$ . Suppose that  $\alpha, \beta \in \mathbb{C}$ . Show that the Laurent series for the function  $\alpha f + \beta g$  has coefficients  $\alpha a_n + \beta b_n$ .



Some more useful exercises:

**Exercise:** Suppose  $f$  and  $g$  are holomorphic functions defined on  $\text{ann}(p; r_1, r_2)$ . Let  $a_n$  be the coefficients in the Laurent series for  $f$  and  $b_n$  be the coefficients in the Laurent series for  $g$ . Suppose that  $\alpha, \beta \in \mathbb{C}$ . Show that the Laurent series for the function  $\alpha f + \beta g$  has coefficients  $\alpha a_n + \beta b_n$ .

**Exercise:** Expand the function  $f(z) = \frac{1}{(z-1)(z-2)}$  in the sets  $\text{ann}(0; 0, 1)$ ,  $\text{ann}(0; 1, 2)$ , and  $\text{ann}(0; 2, \infty)$ .