

Cultivating Complex Analysis: Convergence of sequences of holomorphic functions (3.4.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

What is the right topology for the set of holomorphic functions?

What is the right topology for the set of holomorphic functions?

Curiously, the right topology is the same as for continuous functions:

Definition

A sequence of functions $f_n: U \rightarrow \mathbb{C}$ converges *uniformly on compact subsets* to $f: U \rightarrow \mathbb{C}$ if $f_n|_K$ converges uniformly to $f|_K$ for every compact $K \subset U$.

What is the right topology for the set of holomorphic functions?

Curiously, the right topology is the same as for continuous functions:

Definition

A sequence of functions $f_n: U \rightarrow \mathbb{C}$ converges *uniformly on compact subsets* to $f: U \rightarrow \mathbb{C}$ if $f_n|_K$ converges uniformly to $f|_K$ for every compact $K \subset U$.

What do we mean by the right topology?

The most natural one that preserves holomorphic functions.

What is the right topology for the set of holomorphic functions?

Curiously, the right topology is the same as for continuous functions:

Definition

A sequence of functions $f_n: U \rightarrow \mathbb{C}$ converges *uniformly on compact subsets* to $f: U \rightarrow \mathbb{C}$ if $f_n|_K$ converges uniformly to $f|_K$ for every compact $K \subset U$.

What do we mean by the right topology?

The most natural one that preserves holomorphic functions.

For example, the uniform topology is the right one for continuous functions:

The uniform limit of continuous functions is continuous.

What is the right topology for the set of holomorphic functions?

Curiously, the right topology is the same as for continuous functions:

Definition

A sequence of functions $f_n: U \rightarrow \mathbb{C}$ converges *uniformly on compact subsets* to $f: U \rightarrow \mathbb{C}$ if $f_n|_K$ converges uniformly to $f|_K$ for every compact $K \subset U$.

What do we mean by the right topology?

The most natural one that preserves holomorphic functions.

For example, the uniform topology is the right one for continuous functions:

The uniform limit of continuous functions is continuous.

But for real differentiable functions, it's not:

What is the right topology for the set of holomorphic functions?

Curiously, the right topology is the same as for continuous functions:

Definition

A sequence of functions $f_n: U \rightarrow \mathbb{C}$ converges *uniformly on compact subsets* to $f: U \rightarrow \mathbb{C}$ if $f_n|_K$ converges uniformly to $f|_K$ for every compact $K \subset U$.

What do we mean by the right topology?

The most natural one that preserves holomorphic functions.

For example, the uniform topology is the right one for continuous functions:

The uniform limit of continuous functions is continuous.

But for real differentiable functions, it's not:

$|x|^{1+1/n}$ is C^1 on \mathbb{R} and converges uniformly on compact subsets to $|x|$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

$\partial \Delta_r(p)$ is compact. Let $\delta > 0$ be the distance of z to $\partial \Delta_r(p)$.

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

$\partial \Delta_r(p)$ is compact. Let $\delta > 0$ be the distance of z to $\partial \Delta_r(p)$.

For $\zeta \in \partial \Delta_r(p)$,

$$\left| \frac{f_n(\zeta)}{\zeta - z} - \frac{f(\zeta)}{\zeta - z} \right| = \frac{|f_n(\zeta) - f(\zeta)|}{|\zeta - z|}$$

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

$\partial \Delta_r(p)$ is compact. Let $\delta > 0$ be the distance of z to $\partial \Delta_r(p)$.

For $\zeta \in \partial \Delta_r(p)$,

$$\left| \frac{f_n(\zeta)}{\zeta - z} - \frac{f(\zeta)}{\zeta - z} \right| = \frac{|f_n(\zeta) - f(\zeta)|}{|\zeta - z|} \leq \frac{1}{\delta} |f_n(\zeta) - f(\zeta)|.$$

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

$\partial\Delta_r(p)$ is compact. Let $\delta > 0$ be the distance of z to $\partial\Delta_r(p)$.

For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{f_n(\zeta)}{\zeta - z} - \frac{f(\zeta)}{\zeta - z} \right| = \frac{|f_n(\zeta) - f(\zeta)|}{|\zeta - z|} \leq \frac{1}{\delta} |f_n(\zeta) - f(\zeta)|.$$

I.e., $\zeta \mapsto \frac{f_n(\zeta)}{\zeta - z}$ converges uniformly to $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$ on $\partial\Delta_r(p)$ (as $f_n \rightarrow f$ uniformly on $\partial\Delta_r(p)$).

Theorem

Suppose $U \subset \mathbb{C}$ is open and $f_n: U \rightarrow \mathbb{C}$ is a sequence of holomorphic functions converging uniformly on compact subsets to $f: U \rightarrow \mathbb{C}$. Then f is holomorphic. Moreover, for all ℓ , the ℓ^{th} derivative $f_n^{(\ell)}$ converges uniformly on compact subsets to $f^{(\ell)}$.

Proof: Fix $p \in U$. Consider $\overline{\Delta_r(p)} \subset U$.

For any $z \in \Delta_r(p)$,

$$f_n(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f_n(\zeta)}{\zeta - z} d\zeta.$$

$\partial\Delta_r(p)$ is compact. Let $\delta > 0$ be the distance of z to $\partial\Delta_r(p)$.

For $\zeta \in \partial\Delta_r(p)$,

$$\left| \frac{f_n(\zeta)}{\zeta - z} - \frac{f(\zeta)}{\zeta - z} \right| = \frac{|f_n(\zeta) - f(\zeta)|}{|\zeta - z|} \leq \frac{1}{\delta} |f_n(\zeta) - f(\zeta)|.$$

I.e., $\zeta \mapsto \frac{f_n(\zeta)}{\zeta - z}$ converges uniformly to $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$ on $\partial\Delta_r(p)$ (as $f_n \rightarrow f$ uniformly on $\partial\Delta_r(p)$).

We can take the limit $n \rightarrow \infty$ underneath the integral.

We get
$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p).$$

We get
$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p).$$

$f|_{\partial\Delta_r(p)}$ is continuous by uniform convergence.

We get $f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta$ for all $z \in \Delta_r(p)$.

$f|_{\partial\Delta_r(p)}$ is continuous by uniform convergence.

$f|_{\Delta_r(p)} = C[f|_{\partial\Delta_r(p)}]$ (the Cauchy transform), which is holomorphic.

We get
$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p).$$

$f|_{\partial\Delta_r(p)}$ is continuous by uniform convergence.

$f|_{\Delta_r(p)} = C[f|_{\partial\Delta_r(p)}]$ (the Cauchy transform), which is holomorphic.

So f is holomorphic.

We get
$$f(z) = \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta \quad \text{for all } z \in \Delta_r(p).$$

$f|_{\partial\Delta_r(p)}$ is continuous by uniform convergence.

$f|_{\Delta_r(p)} = C[f|_{\partial\Delta_r(p)}]$ (the Cauchy transform), which is holomorphic.

So f is holomorphic.

We still need to prove the “Moreover” bit:

That $\{f_n^{(\ell)}\}$ also converges uniformly on compact subsets.

Suppose $K \subset U$ is compact.

Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).


Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.




Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.



Suppose $K \subset U$ is compact.


If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).


Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.


$K \subset K' \subset U$.

K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.


K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.


If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.

$\{f_n\}$ converges uniformly on K' :



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.


If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.

$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .


$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.

$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.

For any $p \in K$, by Cauchy estimates on $f_n - f$ in $\Delta_{d/2}(p)$ (note $\partial \Delta_{d/2}(p) \subset K'$):



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.

K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.


Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.

$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.

For any $p \in K$, by Cauchy estimates on $f_n - f$ in $\Delta_{d/2}(p)$ (note $\partial \Delta_{d/2}(p) \subset K'$):

$$\left| f_n^{(\ell)}(p) - f^{(\ell)}(p) \right|$$



Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.


K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.



$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.

For any $p \in K$, by Cauchy estimates on $f_n - f$ in $\Delta_{d/2}(p)$ (note $\partial \Delta_{d/2}(p) \subset K'$):

$$\left| f_n^{(\ell)}(p) - f^{(\ell)}(p) \right| \leq \frac{\ell! \|f_n - f\|_{\partial \Delta_{d/2}(p)}}{(d/2)^\ell}$$

Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.


K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.



$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.

For any $p \in K$, by Cauchy estimates on $f_n - f$ in $\Delta_{d/2}(p)$ (note $\partial\Delta_{d/2}(p) \subset K'$):

$$\left| f_n^{(\ell)}(p) - f^{(\ell)}(p) \right| \leq \frac{\ell! \|f_n - f\|_{\partial\Delta_{d/2}(p)}}{(d/2)^\ell} \leq \frac{\ell! 2^\ell}{d^\ell} \epsilon.$$

Suppose $K \subset U$ is compact.

If $U \neq \mathbb{C}$, the distance d of K and ∂U is positive (if $U = \mathbb{C}$, take $d > 0$ arbitrary).

Let $K' = \bigcup_{z \in K} \overline{\Delta_{d/2}(z)}$.

$K \subset K' \subset U$.


K' is bounded.

If $p \notin K'$, $\exists q \in K$ such that $|p - q|$ is the distance of K (compact) to p .

$|p - q| > d/2 \quad (p \notin K')$.

Every point in $\Delta_{|p-q|-d/2}(p)$ is further than $d/2$ from K (so not in K'). So K' is closed.

$\Rightarrow K'$ is compact.



$\{f_n\}$ converges uniformly on K' : Given $\epsilon > 0$, $\exists N$ such that $|f_n(z) - f(z)| < \epsilon \quad \forall z \in K', n \geq N$.

For any $p \in K$, by Cauchy estimates on $f_n - f$ in $\Delta_{d/2}(p)$ (note $\partial \Delta_{d/2}(p) \subset K'$):

$$\left| f_n^{(\ell)}(p) - f^{(\ell)}(p) \right| \leq \frac{\ell! \|f_n - f\|_{\partial \Delta_{d/2}(p)}}{(d/2)^\ell} \leq \frac{\ell! 2^\ell}{d^\ell} \epsilon.$$

$\Rightarrow \{f_n^{(\ell)}\}$ converges uniformly to $f^{(\ell)}$ on K .

□

Point is: We can write all derivatives of f as integrals of f .

Point is: We can write all derivatives of f as integrals of f .

Integration is a far nicer operation than differentiation, and for holomorphic functions, we can differentiate by integrating.