

Cultivating Complex Analysis:  
Harmonic functions  
Harnack's inequality and principle (7.2.3–7.2.4)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r}$$



## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr}$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)}$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr}$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .  
Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

For  $z = p + re^{i\theta}$  (using  $f \geq 0$ ),

$$f(z) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

For  $z = p + re^{i\theta}$  (using  $f \geq 0$ ),

$$f(z) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt \leq \frac{S+r}{S-r} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) dt \right)$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

For  $z = p + re^{i\theta}$  (using  $f \geq 0$ ),

$$f(z) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt \leq \frac{S+r}{S-r} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) dt \right) = \frac{S+r}{S-r} f(p).$$

## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

For  $z = p + re^{i\theta}$  (using  $f \geq 0$ ),

$$f(z) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt \leq \frac{S+r}{S-r} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) dt \right) = \frac{S+r}{S-r} f(p).$$

The lower inequality follows in the same way.



## Theorem (Harnack's inequality)

Suppose  $f: \Delta_R(p) \rightarrow \mathbb{R}$  is harmonic and nonnegative, and suppose  $0 < r < R$ . Then

$$\frac{R-r}{R+r}f(p) \leq f(z) \leq \frac{R+r}{R-r}f(p) \quad \text{for all } z \in \overline{\Delta_r(p)}.$$

**Proof:**  $\frac{R+r}{R-r}$  increasing in  $r$  and  $\frac{R-r}{R+r}$  decreasing in  $r \Rightarrow$  enough to prove for  $z \in \partial\Delta_r(p)$ .

Let  $S$  be such that  $0 < r < S < R$ .

$$f(p + re^{i\theta}) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} dt.$$

$$\frac{S-r}{S+r} = \frac{S^2 - r^2}{S^2 + r^2 + 2Sr} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr \cos(\theta - t)} \leq \frac{S^2 - r^2}{S^2 + r^2 - 2Sr} = \frac{S+r}{S-r}.$$

For  $z = p + re^{i\theta}$  (using  $f \geq 0$ ),

$$f(z) = \int_{-\pi}^{\pi} f(p + Se^{it}) P_{r/S}(\theta - t) dt \leq \frac{S+r}{S-r} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Se^{it}) dt \right) = \frac{S+r}{S-r} f(p).$$

The lower inequality follows in the same way.

As  $S < R$  was arbitrary, the theorem follows by taking a limit.



The inequalities are optimal.

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

Consider

$$f(z) = \operatorname{Re} \frac{1+z}{1-z}.$$

(the Poisson kernel except for  $\frac{1}{2\pi}$ ).

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

Consider

$$f(z) = \operatorname{Re} \frac{1+z}{1-z}.$$

(the Poisson kernel except for  $\frac{1}{2\pi}$ ).

So  $f(z) > 0$  on  $\mathbb{D}$ .

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

Consider

$$f(z) = \operatorname{Re} \frac{1+z}{1-z}.$$

(the Poisson kernel except for  $\frac{1}{2\pi}$ ).

So  $f(z) > 0$  on  $\mathbb{D}$ .

$$f(0) = 1.$$

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

Consider

$$f(z) = \operatorname{Re} \frac{1+z}{1-z}.$$

(the Poisson kernel except for  $\frac{1}{2\pi}$ ).

So  $f(z) > 0$  on  $\mathbb{D}$ .

$$f(0) = 1.$$

$z = r$  gets equality on the right hand side.

The inequalities are optimal. In  $\mathbb{D}$ , the theorem says

$$\frac{1-r}{1+r}f(0) \leq f(z) \leq \frac{1+r}{1-r}f(0).$$

Consider

$$f(z) = \operatorname{Re} \frac{1+z}{1-z}.$$

(the Poisson kernel except for  $\frac{1}{2\pi}$ ).

So  $f(z) > 0$  on  $\mathbb{D}$ .

$$f(0) = 1.$$

$z = r$  gets equality on the right hand side.

$z = -r$  gets equality on the left hand side.



## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

## Corollary (Harnack's inequality)

*Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that*

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

*for every harmonic and nonnegative function  $f$  defined on  $U$ .*

**Proof:** WLOG assume that  $K$  is connected:

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

$\exists \Delta_r(z_1), \dots, \Delta_r(z_N)$  that cover  $K$  and  $\Delta_{2r}(z_j) \subset U$  for every  $j$ .

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

$\exists \Delta_r(z_1), \dots, \Delta_r(z_N)$  that cover  $K$  and  $\Delta_{2r}(z_j) \subset U$  for every  $j$ .

Fix  $\zeta, \xi \in K$ .

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

$\exists \Delta_r(z_1), \dots, \Delta_r(z_N)$  that cover  $K$  and  $\Delta_{2r}(z_j) \subset U$  for every  $j$ .

Fix  $\zeta, \xi \in K$ . Assume  $\zeta \in \Delta_r(z_1)$

and  $\xi \in \Delta_r(z_n)$

(for some  $n \leq N$ .)

## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

$\exists \Delta_r(z_1), \dots, \Delta_r(z_N)$  that cover  $K$  and  $\Delta_{2r}(z_j) \subset U$  for every  $j$ .

Fix  $\zeta, \xi \in K$ . Assume  $\zeta \in \Delta_r(z_1)$

and  $\xi \in \Delta_r(z_n)$

(for some  $n \leq N$ .)

Also assume  $\Delta_r(z_j) \cap \Delta_r(z_{j+1}) \neq \emptyset$

for all  $j = 1, \dots, n-1$  ( $K$  connected).



## Corollary (Harnack's inequality)

Suppose  $U \subset \mathbb{C}$  is a domain and  $K \subset U$  is compact. Then there exists a  $C > 0$  such that

$$\sup_{z \in K} f(z) \leq C \inf_{z \in K} f(z)$$

for every harmonic and nonnegative function  $f$  defined on  $U$ .

**Proof:** WLOG assume that  $K$  is connected: (e.g. cover by finitely many closed discs so that  $K$  has finitely many components, then connect with paths as  $U$  is path connected).

Let  $r > 0$  be less than half the distance from  $K$  to  $\partial U$ .

$\exists \Delta_r(z_1), \dots, \Delta_r(z_N)$  that cover  $K$  and  $\Delta_{2r}(z_j) \subset U$  for every  $j$ .


Fix  $\zeta, \xi \in K$ . Assume  $\zeta \in \Delta_r(z_1)$

and  $\xi \in \Delta_r(z_n)$

(for some  $n \leq N$ .)

Also assume  $\Delta_r(z_j) \cap \Delta_r(z_{j+1}) \neq \emptyset$

for all  $j = 1, \dots, n-1$  ( $K$  connected).



Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j$

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \quad \Rightarrow \quad$  if  $w \in \Delta_r(z_j)$

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \quad \Rightarrow \quad$  if  $w \in \Delta_r(z_j) \quad \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .



Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .

Let  $q$  be the midpoint between  $z_j$  and  $z_{j+1} \Rightarrow q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1})$ .

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .

Let  $q$  be the midpoint between  $z_j$  and  $z_{j+1} \Rightarrow q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1})$ .

$\Rightarrow f(z_j) \leq 3f(q) \leq 3(3f(z_{j+1})) = 3^2f(z_{j+1})$ .

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$$\Delta_{2r}(z_j) \subset U \text{ for all } j \quad \Rightarrow \quad \text{if } w \in \Delta_r(z_j) \quad \Rightarrow$$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$$\Rightarrow f(w) \leq 3f(z_j) \text{ and } f(z_j) \leq 3f(w).$$

Follow the discs.

$$\zeta \in \Delta_r(z_1) \quad \Rightarrow \quad f(\zeta) \leq 3f(z_1).$$

$$\text{Let } q \text{ be the midpoint between } z_j \text{ and } z_{j+1} \quad \Rightarrow \quad q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1}).$$

$$\Rightarrow f(z_j) \leq 3f(q) \leq 3(3f(z_{j+1})) = 3^2f(z_{j+1}).$$

$$\xi \in \Delta_r(z_n) \quad \Rightarrow \quad f(z_n) \leq 3f(\xi).$$

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .

Let  $q$  be the midpoint between  $z_j$  and  $z_{j+1} \Rightarrow q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1})$ .

$\Rightarrow f(z_j) \leq 3f(q) \leq 3(3f(z_{j+1})) = 3^2f(z_{j+1})$ .

$\xi \in \Delta_r(z_n) \Rightarrow f(z_n) \leq 3f(\xi)$ .

All in all,  $f(\zeta) \leq 3^{2n}f(\xi) \leq 3^{2N}f(\xi)$ .

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .

Let  $q$  be the midpoint between  $z_j$  and  $z_{j+1} \Rightarrow q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1})$ .

$\Rightarrow f(z_j) \leq 3f(q) \leq 3(3f(z_{j+1})) = 3^2f(z_{j+1})$ .

$\xi \in \Delta_r(z_n) \Rightarrow f(z_n) \leq 3f(\xi)$ .

All in all,  $f(\zeta) \leq 3^{2n}f(\xi) \leq 3^{2N}f(\xi)$ .

$N$  only depends on  $K$ , not on  $\zeta, \xi$ , or  $f$ . As  $\zeta$  and  $\xi$  were arbitrary, the theorem follows.  $\square$

Suppose  $f: U \rightarrow \mathbb{R}$  is harmonic and nonnegative.

$\Delta_{2r}(z_j) \subset U$  for all  $j \Rightarrow$  if  $w \in \Delta_r(z_j) \Rightarrow$

$$\frac{1}{3}f(z_j) = \frac{2r-r}{2r+r}f(z_j) \leq f(w) \leq \frac{2r+r}{2r-r}f(z_j) = 3f(z_j).$$

$\Rightarrow f(w) \leq 3f(z_j)$  and  $f(z_j) \leq 3f(w)$ .

Follow the discs.

$\zeta \in \Delta_r(z_1) \Rightarrow f(\zeta) \leq 3f(z_1)$ .

Let  $q$  be the midpoint between  $z_j$  and  $z_{j+1} \Rightarrow q \in \Delta_r(z_j) \cap \Delta_r(z_{j+1})$ .

$\Rightarrow f(z_j) \leq 3f(q) \leq 3(3f(z_{j+1})) = 3^2f(z_{j+1})$ .

$\xi \in \Delta_r(z_n) \Rightarrow f(z_n) \leq 3f(\xi)$ .

All in all,  $f(\zeta) \leq 3^{2n}f(\xi) \leq 3^{2N}f(\xi)$ .

$N$  only depends on  $K$ , not on  $\zeta$ ,  $\xi$ , or  $f$ . As  $\zeta$  and  $\xi$  were arbitrary, the theorem follows.  $\square$

**Remark:** We got an explicit (if not optimal)  $C$ .

**Exercise:** Show by example that Harnack's general inequality need not hold if  $U$  is not assumed to be connected.

**Exercise:** Show by example that Harnack's general inequality need not hold if  $U$  is not assumed to be connected.

**Exercise:** Find the following counterexample of Harnack's inequality if  $f$  is not assumed to be nonnegative. For every  $M > 0$  find a harmonic function  $f: \mathbb{D} \rightarrow \mathbb{R}$  such that  $f(0) = 1$  and  $f(1/2) \geq M$ .



**Exercise:** Show by example that Harnack's general inequality need not hold if  $U$  is not assumed to be connected.

**Exercise:** Find the following counterexample of Harnack's inequality if  $f$  is not assumed to be nonnegative. For every  $M > 0$  find a harmonic function  $f: \mathbb{D} \rightarrow \mathbb{R}$  such that  $f(0) = 1$  and  $f(1/2) \geq M$ .

**Exercise:** Use Harnack's inequality to prove Liouville's theorem for harmonic functions: If  $f: \mathbb{C} \rightarrow \mathbb{R}$  is harmonic and nonnegative, then  $f$  is constant.

## Theorem (Harnack's principle)

*Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.*

## Theorem (Harnack's principle)

*Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.*

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

## Theorem (Harnack's principle)

*Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.*

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).  
By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

## Theorem (Harnack's principle)

*Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.*

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ ,

## Theorem (Harnack's principle)

*Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.*

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

Harnack's inequality  $\Rightarrow f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z)$ .

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

$$\text{Harnack's inequality} \quad \Rightarrow \quad f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z).$$

$$\Rightarrow f_n(z) \rightarrow +\infty \text{ uniformly for } z \in K.$$



## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

$$\text{Harnack's inequality} \quad \Rightarrow \quad f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z).$$

$$\Rightarrow f_n(z) \rightarrow +\infty \text{ uniformly for } z \in K.$$

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

Harnack's inequality  $\Rightarrow f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z)$ .

$\Rightarrow f_n(z) \rightarrow +\infty$  uniformly for  $z \in K$ .

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

Harnack's inequality  $\Rightarrow f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z)$ .

$\Rightarrow f_n(z) \rightarrow +\infty$  uniformly for  $z \in K$ .

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ ,

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

$$\text{Harnack's inequality} \quad \Rightarrow \quad f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z).$$

$$\Rightarrow f_n(z) \rightarrow +\infty \text{ uniformly for } z \in K.$$

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ , then

$$\sup_{z \in K} (f_m(z) - f_n(z)) \leq C \inf_{z \in K} (f_m(z) - f_n(z))$$

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

$$\text{Harnack's inequality} \quad \Rightarrow \quad f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z).$$

$$\Rightarrow f_n(z) \rightarrow +\infty \text{ uniformly for } z \in K.$$

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ , then

$$\sup_{z \in K} (f_m(z) - f_n(z)) \leq C \inf_{z \in K} (f_m(z) - f_n(z)) \leq C(f_m(p) - f_n(p)) < \epsilon.$$

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

$$\text{Harnack's inequality} \quad \Rightarrow \quad f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z).$$

$$\Rightarrow f_n(z) \rightarrow +\infty \text{ uniformly for } z \in K.$$

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ , then

$$\sup_{z \in K} (f_m(z) - f_n(z)) \leq C \inf_{z \in K} (f_m(z) - f_n(z)) \leq C(f_m(p) - f_n(p)) < \epsilon.$$

$$\Rightarrow \{f_n\} \text{ is uniformly Cauchy on } K$$

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

Harnack's inequality  $\Rightarrow f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z)$ .

$\Rightarrow f_n(z) \rightarrow +\infty$  uniformly for  $z \in K$ .

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ , then

$$\sup_{z \in K} (f_m(z) - f_n(z)) \leq C \inf_{z \in K} (f_m(z) - f_n(z)) \leq C(f_m(p) - f_n(p)) < \epsilon.$$

$\Rightarrow \{f_n\}$  is uniformly Cauchy on  $K \Rightarrow$  converges uniformly.

## Theorem (Harnack's principle)

Let  $U \subset \mathbb{C}$  be a domain and  $\{f_n\}$  a sequence of harmonic functions on  $U$  such that  $f_1 \leq f_2 \leq f_3 \leq \dots$ . Then either  $f_n \rightarrow +\infty$  uniformly on compact subsets, or  $f_n \rightarrow f$  for a harmonic  $f: U \rightarrow \mathbb{R}$  uniformly on compact subsets.

**Proof:** WLOG assume  $f_n \geq 0$  for all  $n$ . (Otherwise apply to  $f_n - f_1$ ).

By the monotonicity,  $\{f_n\}$  converges pointwise (possibly to  $+\infty$ ).

If  $\lim f_n(p) = +\infty$  for some  $p$ , let  $K \subset U$  be any compact and let  $K' = K \cup \{p\}$ .

Harnack's inequality  $\Rightarrow f_n(p) \leq \sup_{z \in K'} f_n(z) \leq C \inf_{z \in K'} f_n(z) \leq C \inf_{z \in K} f_n(z)$ .

$\Rightarrow f_n(z) \rightarrow +\infty$  uniformly for  $z \in K$ .

Suppose  $f(z) = \lim f_n(z) < +\infty$  for every  $z \in U$ .

Let  $K \subset U$  be compact, take the  $C$  from Harnack's, and take any  $p \in K$ .

Given  $\epsilon > 0$ , suppose  $m > n$  are such that  $f_m(p) - f_n(p) < \epsilon/C$ , then

$$\sup_{z \in K} (f_m(z) - f_n(z)) \leq C \inf_{z \in K} (f_m(z) - f_n(z)) \leq C(f_m(p) - f_n(p)) < \epsilon.$$

$\Rightarrow \{f_n\}$  is uniformly Cauchy on  $K \Rightarrow$  converges uniformly.

$f$  is harmonic by Harnack's first theorem.





**Exercise:** Prove yet another version of Harnack's principle. Suppose  $U \subset \mathbb{C}$  is a domain,  $\{f_n\}$  is a sequence of nonnegative harmonic functions on  $U$ , and  $p \in U$  is fixed.

- a) If  $f_n(p) \rightarrow +\infty$ , then  $\{f_n\}$  converges to  $+\infty$  uniformly on compact subsets.
- b) If  $f: U \rightarrow \mathbb{R}$  is harmonic,  $f_n(z) \leq f(z)$  for all  $z \in U$ , and  $f_n(p) \rightarrow f(p)$ , then  $\{f_n\}$  converges to  $f$  uniformly on compact subsets.

**Exercise:** Prove yet another version of Harnack's principle. Suppose  $U \subset \mathbb{C}$  is a domain,  $\{f_n\}$  is a sequence of nonnegative harmonic functions on  $U$ , and  $p \in U$  is fixed.

- a) If  $f_n(p) \rightarrow +\infty$ , then  $\{f_n\}$  converges to  $+\infty$  uniformly on compact subsets.
- b) If  $f: U \rightarrow \mathbb{R}$  is harmonic,  $f_n(z) \leq f(z)$  for all  $z \in U$ , and  $f_n(p) \rightarrow f(p)$ , then  $\{f_n\}$  converges to  $f$  uniformly on compact subsets.

**Exercise:** Prove a Montel-like theorem for harmonic functions. Suppose  $U \subset \mathbb{C}$  is a domain and  $\{f_n\}$  is a sequence of nonnegative harmonic functions. Show that at least one (or both) of the following are true:

- (i)  $\exists$  a subsequence converging to  $+\infty$  uniformly on compact subsets.
- (ii)  $\exists$  a subsequence converging to a harmonic function uniformly on compact subsets.