

Cultivating Complex Analysis: Rouché's theorem (5.4.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

Example: z^2 and $(z - \epsilon)(z + \epsilon)$ are close on $\partial\mathbb{D}$.

Two holomorphic functions that are close on a cycle have the same number of zeros inside.

For meromorphic functions the difference of zeros and poles is the same.

Example: z^2 and $(z - \epsilon)(z + \epsilon)$ are close on $\partial\mathbb{D}$.

Example: 1 and $\frac{z-\epsilon}{z+\epsilon}$ are close on $\partial\mathbb{D}$.

Theorem (Rouché)

Suppose $U \subset \mathbb{C}$ is open, Γ is a cycle in U homologous to zero in U , and $n(\Gamma; z)$ is either 0 or 1 for all $z \notin \Gamma$.

Theorem (Rouché)

Suppose $U \subset \mathbb{C}$ is open, Γ is a cycle in U homologous to zero in U , and $n(\Gamma; z)$ is either 0 or 1 for all $z \notin \Gamma$. Suppose that $f: U \rightarrow \mathbb{C}_\infty$ and $g: U \rightarrow \mathbb{C}_\infty$ are meromorphic functions with no zeros or poles on Γ such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Theorem (Rouché)

Suppose $U \subset \mathbb{C}$ is open, Γ is a cycle in U homologous to zero in U , and $n(\Gamma; z)$ is either 0 or 1 for all $z \notin \Gamma$. Suppose that $f: U \rightarrow \mathbb{C}_\infty$ and $g: U \rightarrow \mathbb{C}_\infty$ are meromorphic functions with no zeros or poles on Γ such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$. Let N_f, N_g be the number of zeros in V and P_f, P_g the number of poles in V (both counting multiplicity) of f and g respectively.

Theorem (Rouché)

Suppose $U \subset \mathbb{C}$ is open, Γ is a cycle in U homologous to zero in U , and $n(\Gamma; z)$ is either 0 or 1 for all $z \notin \Gamma$. Suppose that $f: U \rightarrow \mathbb{C}_\infty$ and $g: U \rightarrow \mathbb{C}_\infty$ are meromorphic functions with no zeros or poles on Γ such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$. Let N_f, N_g be the number of zeros in V and P_f, P_g the number of poles in V (both counting multiplicity) of f and g respectively. Then

$$N_f - P_f = N_g - P_g.$$

Theorem (Rouché)

Suppose $U \subset \mathbb{C}$ is open, Γ is a cycle in U homologous to zero in U , and $n(\Gamma; z)$ is either 0 or 1 for all $z \notin \Gamma$. Suppose that $f: U \rightarrow \mathbb{C}_\infty$ and $g: U \rightarrow \mathbb{C}_\infty$ are meromorphic functions with no zeros or poles on Γ such that

$$|f(z) - g(z)| < |f(z)| + |g(z)| \quad \text{for all } z \in \Gamma.$$

Let $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$. Let N_f, N_g be the number of zeros in V and P_f, P_g the number of poles in V (both counting multiplicity) of f and g respectively. Then

$$N_f - P_f = N_g - P_g.$$

Corollary (Rouché)

Let U, Γ and V be as in the theorem. Suppose $f: U \rightarrow \mathbb{C}$ and $g: U \rightarrow \mathbb{C}$ are holomorphic such that $|f(z) - g(z)| < |f(z)| + |g(z)|$ for all $z \in \Gamma$. Then f and g have the same number of zeros (counting multiplicity) in V .


Classical statement of the theorem uses the weaker inequality

$$|f(z) - g(z)| < |f(z)|$$

Classical statement of the theorem uses the weaker inequality

$$|f(z) - g(z)| < |f(z)|$$

It has a nice geometric interpretation:



Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

The function $\frac{\varphi'}{\varphi}$ has an antiderivative $\text{Log} \circ \varphi$ on a neighborhood of Γ .

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

The function $\frac{\varphi'}{\varphi}$ has an antiderivative $\text{Log} \circ \varphi$ on a neighborhood of Γ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$0 = \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz$$

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

The function $\frac{\varphi'}{\varphi}$ has an antiderivative $\text{Log} \circ \varphi$ on a neighborhood of Γ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$0 = \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left(\frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz$$

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

The function $\frac{\varphi'}{\varphi}$ has an antiderivative $\text{Log} \circ \varphi$ on a neighborhood of Γ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left(\frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz - \frac{1}{2\pi i} \int_{\Gamma} \frac{g'(z)}{g(z)} dz \end{aligned}$$

Proof:

$$\left| \frac{f(z)}{g(z)} - 1 \right| < \left| \frac{f(z)}{g(z)} \right| + 1 \quad \text{on } \Gamma.$$

So $\varphi(z) = \frac{f(z)}{g(z)}$ is never negative (and it is never zero) on Γ (hence on a neighborhood).

Let Log be the principal branch of \log on $\mathbb{C} \setminus (-\infty, 0]$.

The function $\frac{\varphi'}{\varphi}$ has an antiderivative $\text{Log} \circ \varphi$ on a neighborhood of Γ .

By Cauchy's theorem for derivatives, together with the argument principle:

$$\begin{aligned} 0 &= \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi'(z)}{\varphi(z)} dz = \frac{1}{2\pi i} \int_{\Gamma} \left(\frac{f'(z)}{f(z)} - \frac{g'(z)}{g(z)} \right) dz \\ &= \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz - \frac{1}{2\pi i} \int_{\Gamma} \frac{g'(z)}{g(z)} dz = (N_f - P_f) - (N_g - P_g). \end{aligned}$$

□

The classical hypothesis $|f(z) - g(z)| < |f(z)|$ is often sufficient.

The classical hypothesis $|f(z) - g(z)| < |f(z)|$ is often sufficient.

Example: Consider $P(z) = z^n + 1$. Let us use Rouché to show that all the zeros are on $\partial\mathbb{D}$.

The classical hypothesis $|f(z) - g(z)| < |f(z)|$ is often sufficient.

Example: Consider $P(z) = z^n + 1$. Let us use Rouché to show that all the zeros are on $\partial\mathbb{D}$.

On $\partial\Delta_{1-\epsilon}(0)$,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

The classical hypothesis $|f(z) - g(z)| < |f(z)|$ is often sufficient.

Example: Consider $P(z) = z^n + 1$. Let us use Rouché to show that all the zeros are on $\partial\mathbb{D}$.

On $\partial\Delta_{1-\epsilon}(0)$,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

By Rouché $P(z)$ and 1 have the same number of zeros in $\Delta_{1-\epsilon}(0)$.

The classical hypothesis $|f(z) - g(z)| < |f(z)|$ is often sufficient.

Example: Consider $P(z) = z^n + 1$. Let us use Rouché to show that all the zeros are on $\partial\mathbb{D}$.

On $\partial\Delta_{1-\epsilon}(0)$,

$$|P(z) - 1| = |z|^n < 1 = |1|.$$

By Rouché $P(z)$ and 1 have the same number of zeros in $\Delta_{1-\epsilon}(0)$.

On $\partial\Delta_{1+\epsilon}(0)$,

$$|P(z) - z^n| = 1 < |z^n|.$$

By Rouché, $P(z)$ and z^n have the same number of zeros in $\Delta_{1+\epsilon}(0)$.

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$.

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$. On $\partial\mathbb{D}$,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$. On $\partial\mathbb{D}$,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$ has zeros at $\pm\sqrt{24}i$ (outside \mathbb{D}) and two zeros at the origin (inside \mathbb{D}). So $P(z)$ also has two zeros in \mathbb{D} .

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$. On $\partial\mathbb{D}$,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$ has zeros at $\pm\sqrt{24}i$ (outside \mathbb{D}) and two zeros at the origin (inside \mathbb{D}). So $P(z)$ also has two zeros in \mathbb{D} .

If $|z| = 46 + \epsilon$, then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$. On $\partial\mathbb{D}$,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$ has zeros at $\pm\sqrt{24}i$ (outside \mathbb{D}) and two zeros at the origin (inside \mathbb{D}). So $P(z)$ also has two zeros in \mathbb{D} .

If $|z| = 46 + \epsilon$, then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

So all four zeros satisfy $|z| < 46 + \epsilon$, that is, $|z| \leq 46$.

Example: Consider $P(z) = z^4 + 12z^3 + 24z^2 + 4z + 6$. On $\partial\mathbb{D}$,

$$\begin{aligned} |P(z) - (z^4 + 24z^2)| &= |12z^3 + 4z + 6| \leq |12z^3| + |4z| + |6| \\ &= 22 < 23 = ||24z^2| - |z^4|| \leq |z^4 + 24z^2|. \end{aligned}$$

$z^4 + 24z^2$ has zeros at $\pm\sqrt{24}i$ (outside \mathbb{D}) and two zeros at the origin (inside \mathbb{D}). So $P(z)$ also has two zeros in \mathbb{D} .

If $|z| = 46 + \epsilon$, then

$$|P(z) - z^4| = |12z^3 + 24z^2 + 4z + 6| \leq 46|z|^3 < |z|^4 = |z^4|.$$

So all four zeros satisfy $|z| < 46 + \epsilon$, that is, $|z| \leq 46$.

(Actually the largest zero of P has modulus less than 10).