

Cultivating Complex Analysis: Cauchy estimates, Liouville, and the fundamental theorem of algebra (3.3.4)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We have the tools to make three fundamental results pop out with little work:

We have the tools to make three fundamental results pop out with little work:

1. The “triangle inequality” on the integral formula for the coefficients of the power series gives estimates on their size: **Cauchy’s estimates**.

We have the tools to make three fundamental results pop out with little work:

1. The “triangle inequality” on the integral formula for the coefficients of the power series gives estimates on their size: **Cauchy’s estimates**.
2. Cauchy’s estimates imply **Liouville’s theorem**: Bounded entire (defined on all of \mathbb{C}) holomorphic functions are constant.

We have the tools to make three fundamental results pop out with little work:

1. The “triangle inequality” on the integral formula for the coefficients of the power series gives estimates on their size: **Cauchy’s estimates**.
2. Cauchy’s estimates imply **Liouville’s theorem**: Bounded entire (defined on all of \mathbb{C}) holomorphic functions are constant.
3. Liouville’s theorem gives **the fundamental theorem of algebra**: Every nonconstant polynomial has a root.

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z - p)^n$.

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z-p)^n$. Then for all n ,

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial \Delta_r(p)}}{r^n}.$$

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z-p)^n$. Then for all n ,

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}.$$

In other words, the sequence $\{|c_n|r^n\}$ is bounded by $\|f\|_{\partial\Delta_r(p)}$.

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z-p)^n$. Then for all n ,

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}.$$

In other words, the sequence $\{|c_n|r^n\}$ is bounded by $\|f\|_{\partial\Delta_r(p)}$.

Proof:

$$|c_n| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta-p)^{n+1}} d\zeta \right|$$

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z-p)^n$. Then for all n ,

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}.$$

In other words, the sequence $\{|c_n|r^n\}$ is bounded by $\|f\|_{\partial\Delta_r(p)}$.

Proof:

$$|c_n| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta-p)^{n+1}} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{\|f\|_{\partial\Delta_r(p)}}{r^{n+1}} |d\zeta|$$

For a set K , denote the *supremum norm* or *uniform norm* of a function:

$$\|f\|_K \stackrel{\text{def}}{=} \sup_{z \in K} |f(z)|.$$

Theorem (Cauchy estimates)

Let $U \subset \mathbb{C}$ be open, $f: U \rightarrow \mathbb{C}$ be holomorphic, and $\overline{\Delta_r(p)} \subset U$ be a closed disc. Expand $f(z) = \sum c_n(z-p)^n$. Then for all n ,

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}.$$

In other words, the sequence $\{|c_n|r^n\}$ is bounded by $\|f\|_{\partial\Delta_r(p)}$.

Proof:

$$|c_n| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta-p)^{n+1}} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{\|f\|_{\partial\Delta_r(p)}}{r^{n+1}} |d\zeta| = \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}. \quad \square$$

A better estimate than

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}$$

is not possible.

A better estimate than

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}$$

is not possible.

Consider

$$f(z) = \frac{M}{r^n} (z - p)^n.$$

A better estimate than

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}$$

is not possible.

Consider

$$f(z) = \frac{M}{r^n} (z - p)^n.$$

Then

$$M = \|f\|_{\partial\Delta_r(p)}.$$

A better estimate than

$$|c_n| = \left| \frac{f^{(n)}(p)}{n!} \right| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n}$$

is not possible.

Consider

$$f(z) = \frac{M}{r^n} (z - p)^n.$$

Then

$$M = \|f\|_{\partial\Delta_r(p)}.$$

And

$$|c_n| = \frac{M}{r^n}.$$

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$.

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Expand around the origin:

$$f(z) = \sum_{n=0}^{\infty} c_n z^n.$$

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Expand around the origin:

$$f(z) = \sum_{n=0}^{\infty} c_n z^n.$$

f is holomorphic on a disc of arbitrary radius

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Expand around the origin:

$$f(z) = \sum_{n=0}^{\infty} c_n z^n.$$

f is holomorphic on a disc of arbitrary radius \Rightarrow the Cauchy estimates say

$$|c_n| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n} \leq \frac{M}{r^n} \quad \text{for all } r > 0.$$

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Expand around the origin:

$$f(z) = \sum_{n=0}^{\infty} c_n z^n.$$

f is holomorphic on a disc of arbitrary radius \Rightarrow the Cauchy estimates say

$$|c_n| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n} \leq \frac{M}{r^n} \quad \text{for all } r > 0.$$

Letting $r \rightarrow \infty$ shows that $c_n = 0$ for $n \geq 1$.

Definition

A holomorphic $f: \mathbb{C} \rightarrow \mathbb{C}$ is called an *entire holomorphic function* or perhaps just *entire*.

Examples: Polynomials, e^z , $\sin z$, $\cos z$, but not $\frac{1}{z}$.

Theorem (Liouville)

If f is entire and bounded, then f is constant.

Proof: Suppose f is entire and $|f(z)| \leq M$ for all $z \in \mathbb{C}$. Expand around the origin:

$$f(z) = \sum_{n=0}^{\infty} c_n z^n.$$

f is holomorphic on a disc of arbitrary radius \Rightarrow the Cauchy estimates say

$$|c_n| \leq \frac{\|f\|_{\partial\Delta_r(p)}}{r^n} \leq \frac{M}{r^n} \quad \text{for all } r > 0.$$

Letting $r \rightarrow \infty$ shows that $c_n = 0$ for $n \geq 1$. In other words, $f(z) = c_0$ for all z .



Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Proof: If $P(z)$ does not have a root, then $R(z) = \frac{1}{P(z)}$ is an entire holomorphic function.

Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Proof: If $P(z)$ does not have a root, then $R(z) = \frac{1}{P(z)}$ is an entire holomorphic function.

Suppose $P(z)$ is nonconstant. Then (via an exercise)

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Proof: If $P(z)$ does not have a root, then $R(z) = \frac{1}{P(z)}$ is an entire holomorphic function.

Suppose $P(z)$ is nonconstant. Then (via an exercise)

$$\lim_{z \rightarrow \infty} P(z) = \infty \quad \Rightarrow \quad \lim_{z \rightarrow \infty} R(z) = 0.$$

Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Proof: If $P(z)$ does not have a root, then $R(z) = \frac{1}{P(z)}$ is an entire holomorphic function.

Suppose $P(z)$ is nonconstant. Then (via an exercise)

$$\lim_{z \rightarrow \infty} P(z) = \infty \quad \Rightarrow \quad \lim_{z \rightarrow \infty} R(z) = 0.$$

So $R(z)$ is bounded.

Theorem (Fundamental theorem of algebra)

If $P(z)$ is a nonconstant polynomial, then P has a root.

Proof: If $P(z)$ does not have a root, then $R(z) = \frac{1}{P(z)}$ is an entire holomorphic function.

Suppose $P(z)$ is nonconstant. Then (via an exercise)

$$\lim_{z \rightarrow \infty} P(z) = \infty \quad \Rightarrow \quad \lim_{z \rightarrow \infty} R(z) = 0.$$

So $R(z)$ is bounded.

Liouville says that $R(z)$ and hence $P(z)$ must be constant, a contradiction.

