

Cultivating Complex Analysis: Simply connected domains (4.3 part 2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).*

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).*
- (ii) Every holomorphic $f: U \rightarrow \mathbb{C}$ has a primitive.*

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).*
- (ii) Every holomorphic $f: U \rightarrow \mathbb{C}$ has a primitive.*
- (iii) For every nowhere zero holomorphic $f: U \rightarrow \mathbb{C}$ there exists a holomorphic $g: U \rightarrow \mathbb{C}$ such that $e^{g(z)} = f(z)$.*

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).
- (ii) Every holomorphic $f: U \rightarrow \mathbb{C}$ has a primitive.
- (iii) For every nowhere zero holomorphic $f: U \rightarrow \mathbb{C}$ there exists a holomorphic $g: U \rightarrow \mathbb{C}$ such that $e^{g(z)} = f(z)$.
- (iv) $\frac{1}{z-p}$ has a primitive in U for every $p \in \mathbb{C} \setminus U$.

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).
- (ii) Every holomorphic $f: U \rightarrow \mathbb{C}$ has a primitive.
- (iii) For every nowhere zero holomorphic $f: U \rightarrow \mathbb{C}$ there exists a holomorphic $g: U \rightarrow \mathbb{C}$ such that $e^{g(z)} = f(z)$.
- (iv) $\frac{1}{z-p}$ has a primitive in U for every $p \in \mathbb{C} \setminus U$.
- (v) For every holomorphic $f: U \rightarrow \mathbb{C}$ and every cycle Γ in U , $\int_{\Gamma} f(z) dz = 0$.

There is a whole list of conditions equivalent to being simply connected for domains in \mathbb{C} .

Proposition

Let $U \subset \mathbb{C}$ be a domain. The following are equivalent:

- (i) U is simply connected (in the homology sense).
- (ii) Every holomorphic $f: U \rightarrow \mathbb{C}$ has a primitive.
- (iii) For every nowhere zero holomorphic $f: U \rightarrow \mathbb{C}$ there exists a holomorphic $g: U \rightarrow \mathbb{C}$ such that $e^{g(z)} = f(z)$.
- (iv) $\frac{1}{z-p}$ has a primitive in U for every $p \in \mathbb{C} \setminus U$.
- (v) For every holomorphic $f: U \rightarrow \mathbb{C}$ and every cycle Γ in U , $\int_{\Gamma} f(z) dz = 0$.
- (vi) For every $p \in \mathbb{C} \setminus U$ and every cycle Γ in U , $\int_{\Gamma} \frac{1}{z-p} dz = 0$.

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p]$$

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} \left[e^{g(z)} \right]$$

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} \left[e^{g(z)} \right] = e^{g(z)} g'(z)$$

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} \left[e^{g(z)} \right] = e^{g(z)} g'(z) = (z - p) g'(z).$$

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} \left[e^{g(z)} \right] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} [e^{g(z)}] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$,

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} [e^{g(z)}] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$, so (vi) holds.

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} [e^{g(z)}] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$, so (vi) holds.

As $n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$, (vi) is a restatement of (i).

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} [e^{g(z)}] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$, so (vi) holds.

As $n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$, (vi) is a restatement of (i).

By Cauchy's theorem for derivatives, (ii) \Rightarrow (v) (Cauchy in U).

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} \left[e^{g(z)} \right] = e^{g(z)} g'(z) = (z - p) g'(z).$$

So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$, so (vi) holds.

As $n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$, (vi) is a restatement of (i).

By Cauchy's theorem for derivatives, (ii) \Rightarrow (v) (Cauchy in U).

(v) \Rightarrow (vi) is immediate.

Proof: We just proved (i) (simply connected) \Rightarrow (ii) (existence of primitives), and then we proved (ii) \Rightarrow (iii) (existence of logs).

Suppose (iii).

For $p \in \mathbb{C} \setminus U$, find a g such that $e^{g(z)} = z - p$.

$$1 = \frac{d}{dz} [z - p] = \frac{d}{dz} [e^{g(z)}] = e^{g(z)} g'(z) = (z - p) g'(z).$$


So (iv) holds ($\frac{1}{z-p}$ has a primitive).

By Cauchy's theorem for derivatives, (iv) implies $\int_{\Gamma} \frac{1}{z-p} dz = 0 \quad \forall p \in \mathbb{C} \setminus U$, so (vi) holds.

As $n(\Gamma; p) = \frac{1}{2\pi i} \int_{\Gamma} \frac{1}{z-p} dz$, (vi) is a restatement of (i).

By Cauchy's theorem for derivatives, (ii) \Rightarrow (v) (Cauchy in U).

(v) \Rightarrow (vi) is immediate.



Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following
(although we'll only prove one direction right now).

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma$

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected)

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected) \Rightarrow so $\varphi|_S$ is constant.

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected) \Rightarrow so $\varphi|_S$ is constant.

$\varphi(\infty) = 0, \infty \in S \Rightarrow \varphi|_S \equiv 0$

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected) \Rightarrow so $\varphi|_S$ is constant.

$\varphi(\infty) = 0, \infty \in S \Rightarrow \varphi|_S \equiv 0 \Rightarrow U$ is simply connected. □

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected) \Rightarrow so $\varphi|_S$ is constant.

$\varphi(\infty) = 0, \infty \in S \Rightarrow \varphi|_S \equiv 0 \Rightarrow U$ is simply connected. □

Remark: It is important to use \mathbb{C}_∞ and not \mathbb{C} :

Remark: The existence of roots can also be on the list, but it will be easier to prove later.

Another criterion that can be put on the list is the following (although we'll only prove one direction right now).

Proposition

Let $U \subset \mathbb{C}$ be a domain. If $\mathbb{C}_\infty \setminus U$ is connected, then U is simply connected.

Proof: Take $S = \mathbb{C}_\infty \setminus U$ and let Γ be a cycle in U .

$\varphi(z) = n(\Gamma; z)$ is continuous on $\mathbb{C} \setminus \Gamma \Rightarrow \varphi$ is continuous on $S \setminus \{\infty\}$.

On the unbounded component of $\mathbb{C} \setminus \Gamma$, we have $\varphi = 0$, so $\varphi = 0$ in a neighborhood of ∞ .

Set $\varphi(\infty) = 0$ to make a continuous function on $\mathbb{C}_\infty \setminus \Gamma$ (hence on S).

S is contained in a single component of $\mathbb{C}_\infty \setminus \Gamma$ (S is connected) \Rightarrow so $\varphi|_S$ is constant.

$\varphi(\infty) = 0, \infty \in S \Rightarrow \varphi|_S \equiv 0 \Rightarrow U$ is simply connected. □

Remark: It is important to use \mathbb{C}_∞ and not \mathbb{C} : If $U = \mathbb{C} \setminus \{0\}$ (not simply connected), then $\mathbb{C} \setminus U = \{0\}$ is connected, but $\mathbb{C}_\infty \setminus U = \{0, \infty\}$ is not connected.

Exercise: Let $U_1, U_2 \subset \mathbb{C}$ be two simply connected domains.

Exercise: Let $U_1, U_2 \subset \mathbb{C}$ be two simply connected domains.

1) $U_1 \cup U_2$ is not necessarily simply connected.

Exercise: Let $U_1, U_2 \subset \mathbb{C}$ be two simply connected domains.

- 1) $U_1 \cup U_2$ is not necessarily simply connected.
- 2) If $U_1 \cap U_2$ is nonempty and connected, then $U_1 \cup U_2$ is simply connected.