

Cultivating Complex Analysis: Schwarz's lemma (3.5.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

- (i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

- (i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and
- (ii) $|f'(0)| \leq 1$.

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

- (i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and
- (ii) $|f'(0)| \leq 1$.

Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$ or $|f'(0)| = 1$, then there is a $\theta \in \mathbb{R}$ such that $f(z) = e^{i\theta} z$ for all $z \in \mathbb{D}$.

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

(i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and

(ii) $|f'(0)| \leq 1$.

Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$ or $|f'(0)| = 1$, then there is a $\theta \in \mathbb{R}$ such that $f(z) = e^{i\theta}z$ for all $z \in \mathbb{D}$.

This may sound very specialized, but

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

(i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and

(ii) $|f'(0)| \leq 1$.

Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$ or $|f'(0)| = 1$, then there is a $\theta \in \mathbb{R}$ such that $f(z) = e^{i\theta} z$ for all $z \in \mathbb{D}$.

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into \mathbb{D} .

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

(i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and

(ii) $|f'(0)| \leq 1$.

Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$ or $|f'(0)| = 1$, then there is a $\theta \in \mathbb{R}$ such that $f(z) = e^{i\theta} z$ for all $z \in \mathbb{D}$.

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into \mathbb{D} .
The lemma is telling us about local behavior of a holomorphic function.

Lemma (Schwarz's lemma)

Suppose $f: \mathbb{D} \rightarrow \mathbb{D}$ is holomorphic and $f(0) = 0$, then

(i) $|f(z)| \leq |z|$ for all $z \in \mathbb{D}$, and

(ii) $|f'(0)| \leq 1$.

Furthermore, if $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$ or $|f'(0)| = 1$, then there is a $\theta \in \mathbb{R}$ such that $f(z) = e^{i\theta}z$ for all $z \in \mathbb{D}$.

This may sound very specialized, but

a) A disc is a basic neighborhood and any disc can be translated and rescaled into \mathbb{D} .

The lemma is telling us about local behavior of a holomorphic function.

b) We'll see later that every domain "without holes" (except \mathbb{C}) is biholomorphic to \mathbb{D} , so it tells us about global behavior as well.

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1}$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|}$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$.

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D},$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\}$

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$ attains a maximum in \mathbb{D}

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$ attains a maximum in $\mathbb{D} \Rightarrow g$ is constant

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = zg(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$ attains a maximum in $\mathbb{D} \Rightarrow g$ is constant $\Rightarrow f(z) = e^{i\theta} z$.

Proof: As $f(0) = 0$,

$$f(z) = \sum_{n=1}^{\infty} c_n z^n = z \sum_{n=1}^{\infty} c_n z^{n-1} = z g(z) \quad (g(z) \text{ a holomorphic function on } \mathbb{D}.)$$

Take $0 < r < 1$. For $z \in \partial\Delta_r(0)$,

$$|g(z)| = \frac{|f(z)|}{|z|} \leq \frac{1}{r}.$$

By the maximum modulus principle, $|g(z)| \leq \frac{1}{r}$ for all $z \in \Delta_r(0)$.

Fix any $z \in \mathbb{D}$ and take the limit as $r \uparrow 1$ to find $|g(z)| \leq 1$. So

$$|f(z)| \leq |z| \quad \text{for all } z \in \mathbb{D}, \quad \text{and} \quad |f'(0)| = \left| \lim_{z \rightarrow 0} \frac{f(z)}{z} \right| = |g(0)| \leq 1.$$

If $|f(z_0)| = |z_0|$ for some $z_0 \in \mathbb{D} \setminus \{0\} \Rightarrow g$ attains a maximum in $\mathbb{D} \Rightarrow g$ is constant $\Rightarrow f(z) = e^{i\theta} z$.

As $g(0) = f'(0)$, the same conclusion holds if $|f'(0)| = 1$.



Consider the statement for $f(z) = z^n$ for $n > 1$.

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

For $z \in \mathbb{D} \setminus \{0\}$,

$$|z^n| = |z|^n < |z|.$$

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

For $z \in \mathbb{D} \setminus \{0\}$,

$$|z^n| = |z|^n < |z|.$$

As $f'(z) = nz^{n-1}$, $|f'(0)| = 0 < 1$.

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

For $z \in \mathbb{D} \setminus \{0\}$,

$$|z^n| = |z|^n < |z|.$$

As $f'(z) = nz^{n-1}$, $|f'(0)| = 0 < 1$.

A bound on the derivative does not hold at other points:
Picking the right z and n , can make $|f'(z)|$ arbitrarily large.

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

For $z \in \mathbb{D} \setminus \{0\}$,

$$|z^n| = |z|^n < |z|.$$

As $f'(z) = nz^{n-1}$, $|f'(0)| = 0 < 1$.

A bound on the derivative does not hold at other points:

Picking the right z and n , can make $|f'(z)|$ arbitrarily large.

We can make $|z^n|$ arbitrarily small for a fixed $z \in \mathbb{D}$ by picking a large enough n , but we cannot make it bigger than $|z|$.

Consider the statement for $f(z) = z^n$ for $n > 1$.

f takes \mathbb{D} to \mathbb{D} and $f(0) = 0$.

For $z \in \mathbb{D} \setminus \{0\}$,

$$|z^n| = |z|^n < |z|.$$

As $f'(z) = nz^{n-1}$, $|f'(0)| = 0 < 1$.

A bound on the derivative does not hold at other points:

Picking the right z and n , can make $|f'(z)|$ arbitrarily large.

We can make $|z^n|$ arbitrarily small for a fixed $z \in \mathbb{D}$ by picking a large enough n , but we cannot make it bigger than $|z|$.

Schwarz's lemma says all holomorphic functions behave this way, not just z^n .

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose $U \subset \mathbb{C}$ is bounded, $f: U \rightarrow U$ is holomorphic, $p \in U$, and $f(p) = p$. Then

- (i) $|f'(p)| \leq 1$.
- (ii) If $f'(p) = 1$, then $f(z) = z$ for all $z \in U$.

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose $U \subset \mathbb{C}$ is bounded, $f: U \rightarrow U$ is holomorphic, $p \in U$, and $f(p) = p$. Then

(i) $|f'(p)| \leq 1$.

(ii) If $f'(p) = 1$, then $f(z) = z$ for all $z \in U$.

Hint: WLOG $p = 0$,

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose $U \subset \mathbb{C}$ is bounded, $f: U \rightarrow U$ is holomorphic, $p \in U$, and $f(p) = p$. Then

(i) $|f'(p)| \leq 1$.

(ii) If $f'(p) = 1$, then $f(z) = z$ for all $z \in U$.

Hint: WLOG $p = 0$, then consider the power series expansions of f^ℓ , the ℓ^{th} composition of f with itself, $f(f(f(\cdots f(z) \cdots)))$.

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose $U \subset \mathbb{C}$ is bounded, $f: U \rightarrow U$ is holomorphic, $p \in U$, and $f(p) = p$. Then

(i) $|f'(p)| \leq 1$.

(ii) If $f'(p) = 1$, then $f(z) = z$ for all $z \in U$.

Hint: WLOG $p = 0$, then consider the power series expansions of f^ℓ , the ℓ^{th} composition of f with itself, $f(f(f(\cdots f(z) \cdots)))$.

For (i) consider the linear term of f^ℓ .

It is a useful exercise to rewrite the lemma for $f: \Delta_r(p) \rightarrow \Delta_s(q)$ with $f(p) = q$.

Using an LFT we can find a half-plane ($\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$) version:

If $f: \mathbb{H} \rightarrow \mathbb{H}$ holomorphic and $f(i) = i$, then

$$\left| \frac{f(z) - i}{\overline{f(z)} - i} \right| \leq \left| \frac{z - i}{\bar{z} - i} \right| \quad \text{and} \quad |f'(i)| \leq 1.$$

Another interesting (and more challenging) exercise is the following generalization (sometimes called Cartan's uniqueness theorem)

Suppose $U \subset \mathbb{C}$ is bounded, $f: U \rightarrow U$ is holomorphic, $p \in U$, and $f(p) = p$. Then

(i) $|f'(p)| \leq 1$.

(ii) If $f'(p) = 1$, then $f(z) = z$ for all $z \in U$.

Hint: WLOG $p = 0$, then consider the power series expansions of f^ℓ , the ℓ^{th} composition of f with itself, $f(f(f(\cdots f(z) \cdots)))$.

For (i) consider the linear term of f^ℓ .

For (ii) use Cauchy estimates on the first nonzero nonlinear term of f^ℓ .