

Cultivating Complex Analysis:
Harmonic functions
Dirichlet problem in a disc and the Poisson kernel (7.2.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Dirichlet problem:

Dirichlet problem:

$U \subset \mathbb{C}$ open,

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous,

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Solvable for many U , but not all.

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Solvable for many U , but not all. But solution unique if it exists:

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Solvable for many U , but not all. But solution unique if it exists:

Proposition

Suppose $U \subset \mathbb{C}$ is a bounded domain, $f, g: \bar{U} \rightarrow \mathbb{R}$ are continuous functions harmonic on U , such that $f = g$ on ∂U . Then $f = g$ on U .

Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Solvable for many U , but not all. But solution unique if it exists:

Proposition

Suppose $U \subset \mathbb{C}$ is a bounded domain, $f, g: \bar{U} \rightarrow \mathbb{R}$ are continuous functions harmonic on U , such that $f = g$ on ∂U . Then $f = g$ on U .

Proof: Maximum principle (second version) on $f - g$.



Dirichlet problem:

$U \subset \mathbb{C}$ open, $f: \partial U \rightarrow \mathbb{R}$ continuous, find a continuous $g: \bar{U} \rightarrow \mathbb{R}$ harmonic on U , such that $g|_{\partial U} = f$.

Solvable for many U , but not all. But solution unique if it exists:

Proposition

Suppose $U \subset \mathbb{C}$ is a bounded domain, $f, g: \bar{U} \rightarrow \mathbb{R}$ are continuous functions harmonic on U , such that $f = g$ on ∂U . Then $f = g$ on U .

Proof: Maximum principle (second version) on $f - g$. □

On an unbounded domain, solution need not be unique.


Exercise: The Dirichlet problem does not have a unique solution on the upper half plane \mathbb{H} .

The Poisson kernel for the unit disc $\mathbb{D} \subset \mathbb{C}$ is

$$P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad \text{for } 0 \leq r < 1.$$

As a function of $z = re^{i\theta}$:

$$\frac{1}{2\pi} \operatorname{Re} \left(\frac{1+z}{1-z} \right)$$



Proposition

(i) $P_r(\theta) > 0$ for all $0 \leq r < 1$ and all θ .

Proposition

- (i) $P_r(\theta) > 0$ for all $0 \leq r < 1$ and all θ .
- (ii) $\int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$ for all $0 \leq r < 1$.


Proposition

- (i) $P_r(\theta) > 0$ for all $0 \leq r < 1$ and all θ .
- (ii) $\int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$ for all $0 \leq r < 1$.
- (iii) For any given $\delta > 0$, $\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0$ as $r \uparrow 1$.

Proposition

- (i) $P_r(\theta) > 0$ for all $0 \leq r < 1$ and all θ .
- (ii) $\int_{-\pi}^{\pi} P_r(\theta) d\theta = 1$ for all $0 \leq r < 1$.
- (iii) For any given $\delta > 0$, $\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0$ as $r \uparrow 1$.

The graph of P_r as a function of θ on $[-\pi, \pi]$ for $r = 0.5$, $r = 0.7$, and $r = 0.85$:



Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\int_{-\pi}^{\pi} P_r(\theta) d\theta$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\int_{-\pi}^{\pi} P_r(\theta) d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1 + re^{i\theta}}{1 - re^{i\theta}} \right) d\theta$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\int_{-\pi}^{\pi} P_r(\theta) d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \end{aligned}$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1):$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$.

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$. On $(0, \pi)$, P_r is strictly decreasing, so only need to show that $P_r(\delta) \rightarrow 0$.

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$. On $(0, \pi)$, P_r is strictly decreasing, so only need to show that $P_r(\delta) \rightarrow 0$. $r \mapsto \frac{1+re^{i\delta}}{1-re^{i\delta}}$ is continuous at $r = 1$ (as $\delta > 0$) and

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$. On $(0, \pi)$, P_r is strictly decreasing, so only need to show that $P_r(\delta) \rightarrow 0$. $r \mapsto \frac{1+re^{i\delta}}{1-re^{i\delta}}$ is continuous at $r = 1$ (as $\delta > 0$) and

$$\frac{1+e^{i\delta}}{1-e^{i\delta}} = \frac{(1+e^{i\delta})(1-e^{-i\delta})}{(1-e^{i\delta})(1-e^{-i\delta})}$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$. On $(0, \pi)$, P_r is strictly decreasing, so only need to show that $P_r(\delta) \rightarrow 0$. $r \mapsto \frac{1+re^{i\delta}}{1-re^{i\delta}}$ is continuous at $r = 1$ (as $\delta > 0$) and

$$\frac{1+e^{i\delta}}{1-e^{i\delta}} = \frac{(1+e^{i\delta})(1-e^{-i\delta})}{(1-e^{i\delta})(1-e^{-i\delta})} = \frac{e^{i\delta} - e^{-i\delta}}{|1-e^{i\delta}|^2}$$

Recall: $P_r(\theta) = \frac{1}{2\pi} \frac{1-r^2}{1+r^2-2r\cos\theta} = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right), \quad 0 \leq r < 1.$

(i) (positive): $1 - r^2 > 0$ and $1 + r^2 - 2r \cos \theta \geq 1 + r^2 - 2r = (1 - r)^2 > 0$

(ii) (integral is 1):

$$\begin{aligned} \int_{-\pi}^{\pi} P_r(\theta) d\theta &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \operatorname{Re} \left(\frac{1+re^{i\theta}}{1-re^{i\theta}} \right) d\theta = \operatorname{Re} \frac{1}{2\pi i} \int_{-\pi}^{\pi} \frac{1+re^{i\theta}}{1-re^{i\theta}} \frac{1}{re^{i\theta}} ire^{i\theta} d\theta \\ &= \operatorname{Re} \frac{1}{2\pi i} \int_{\partial\Delta_r(0)} \frac{(1+z)/(1-z)}{z} dz \stackrel{\text{CIF}}{=} \operatorname{Re} \frac{1+0}{1-0} = 1. \end{aligned}$$

(iii) $(\sup\{P_r(\theta) : \delta \leq |\theta| \leq \pi\} \rightarrow 0 \text{ as } r \uparrow 1)$:

By symmetry only need to prove for $\delta \leq \theta \leq \pi$. On $(0, \pi)$, P_r is strictly decreasing, so only need to show that $P_r(\delta) \rightarrow 0$. $r \mapsto \frac{1+re^{i\delta}}{1-re^{i\delta}}$ is continuous at $r = 1$ (as $\delta > 0$) and

$$\frac{1+e^{i\delta}}{1-e^{i\delta}} = \frac{(1+e^{i\delta})(1-e^{-i\delta})}{(1-e^{i\delta})(1-e^{-i\delta})} = \frac{e^{i\delta} - e^{-i\delta}}{|1-e^{i\delta}|^2} = i \frac{2 \operatorname{Im} e^{i\delta}}{|1-e^{i\delta}|^2}. \quad \square$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$.

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t)$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right)$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

is harmonic as a function of $z = re^{i\theta}$.

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

is harmonic as a function of $z = re^{i\theta}$.

$$Pf(z) = Pf(re^{i\theta})$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

is harmonic as a function of $z = re^{i\theta}$.

$$Pf(z) = Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

is harmonic as a function of $z = re^{i\theta}$.

$$Pf(z) = Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right) dt$$

Theorem

Let $f: \partial\mathbb{D} \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\mathbb{D}} \rightarrow \mathbb{R}$, defined by

$$Pf(re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt & \text{if } r < 1, \\ f(e^{i\theta}) & \text{if } r = 1, \end{cases}$$

is harmonic in \mathbb{D} and continuous on $\overline{\mathbb{D}}$.

Proof: Let $z = re^{i\theta}$. For fixed t ,

$$P_r(\theta - t) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + re^{i(\theta-t)}}{1 - re^{i(\theta-t)}} \right) = \frac{1}{2\pi} \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right)$$

is harmonic as a function of $z = re^{i\theta}$.

$$Pf(z) = Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) \operatorname{Re} \left(\frac{1 + ze^{-it}}{1 - ze^{-it}} \right) dt$$

is harmonic for $z = re^{i\theta} \in \mathbb{D}$ (differentiate under the integral).

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given.

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}.$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}.$

f uniformly continuous

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}.$

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta.$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})|$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right|$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

First:

$$\left| \int_{-\pi}^{-\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right|$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

First:

$$\left| \int_{-\pi}^{-\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \int_{-\pi}^{-\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

First:

$$\left| \int_{-\pi}^{-\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \int_{-\pi}^{-\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq (\pi - \delta) 2M \frac{\epsilon}{8M\pi}$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

First:

$$\left| \int_{-\pi}^{-\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \int_{-\pi}^{-\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq (\pi - \delta)2M \frac{\epsilon}{8M\pi} < \frac{\epsilon}{4}.$$

$$Pf(re^{i\theta}) = \int_{-\pi}^{\pi} f(e^{it})P_r(\theta - t) dt = \int_{-\pi}^{\pi} f(e^{i(\theta-t)})P_r(t) dt. \quad (2\pi\text{-periodic})$$

Let $\epsilon > 0$ be given. $M = \sup\{|f(z)| : z \in \partial\mathbb{D}\}$.

f uniformly continuous $\Rightarrow \exists \delta > 0$ such that $|f(e^{i(\theta-t)}) - f(e^{i\theta})| < \epsilon/2$ whenever $|t| < \delta$.

Find $\delta' > 0$ such that $0 < P_r(t) < \frac{\epsilon}{8M\pi}$ whenever $1 - \delta' < r < 1$ and $|t| \geq \delta$.

$$f(e^{i\theta}) = \int_{-\pi}^{\pi} f(e^{i\theta})P_r(t) dt \quad \left(\text{as } \int_{-\pi}^{\pi} P_r(t) dt = 1 \right).$$

$$|Pf(re^{i\theta}) - f(e^{i\theta})| = \left| \int_{-\pi}^{\pi} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \left| \int_{-\pi}^{-\delta} \cdots dt \right| + \left| \int_{-\delta}^{\delta} \cdots dt \right| + \left| \int_{\delta}^{\pi} \cdots dt \right|.$$

First:

$$\left| \int_{-\pi}^{-\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta}))P_r(t) dt \right| \leq \int_{-\pi}^{-\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq (\pi - \delta)2M \frac{\epsilon}{8M\pi} < \frac{\epsilon}{4}.$$

The integral from δ to π is exactly the same.

$$\left|\int_{-\delta}^{\delta}\left(f\left(e^{i(\theta-t)}\right)-f\left(e^{i \theta}\right)\right) P_r(t) d t\right|$$

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt$$

$$\begin{aligned} \left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| &\leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt \\ &\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt \end{aligned}$$

$$\begin{aligned}
 \left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| &\leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt \\
 &\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.
 \end{aligned}$$

$$\begin{aligned}
\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| &\leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt \\
&\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.
\end{aligned}$$

So $1 - \delta' < r < 1 \quad \Rightarrow \quad |Pf(re^{i\theta}) - f(e^{i\theta})|$

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \quad \Rightarrow \quad |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \implies |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$\implies Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \implies |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$\implies Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$.

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that

$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$

$$\left| \int_{-\delta}^{\delta} \left(f(e^{i(\theta-t)}) - f(e^{i\theta}) \right) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} \left| f(e^{i(\theta-t)}) - f(e^{i\theta}) \right| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that

$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$

For possibly smaller δ

$$1 - \delta < r \leq 1 \Rightarrow |Pf(re^{i\theta}) - Pf(e^{i\theta})| < \epsilon/2 \quad \text{for all } \theta.$$

$$\begin{aligned} \left| \int_{-\delta}^{\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta})) P_r(t) dt \right| &\leq \int_{-\delta}^{\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt \\ &\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}. \end{aligned}$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$

$\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that

$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$

For possibly smaller δ

$$1 - \delta < r \leq 1 \Rightarrow |Pf(re^{i\theta}) - Pf(e^{i\theta})| < \epsilon/2 \quad \text{for all } \theta.$$

So if $z = re^{i\theta}$ satisfies $1 - \delta < r \leq 1$ and $|\theta - \theta_0| < \delta$,

$$\left| \int_{-\delta}^{\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta})) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$
 $\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that


$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$

For possibly smaller δ

$$1 - \delta < r \leq 1 \Rightarrow |Pf(re^{i\theta}) - Pf(e^{i\theta})| < \epsilon/2 \quad \text{for all } \theta.$$

So if $z = re^{i\theta}$ satisfies $1 - \delta < r \leq 1$ and $|\theta - \theta_0| < \delta$,

$$|Pf(re^{i\theta}) - Pf(e^{i\theta_0})| \leq |Pf(re^{i\theta}) - Pf(e^{i\theta})| + |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon.$$



$$\left| \int_{-\delta}^{\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta})) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$
 $\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that

$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$


For possibly smaller δ

$$1 - \delta < r \leq 1 \Rightarrow |Pf(re^{i\theta}) - Pf(e^{i\theta})| < \epsilon/2 \quad \text{for all } \theta.$$

So if $z = re^{i\theta}$ satisfies $1 - \delta < r \leq 1$ and $|\theta - \theta_0| < \delta$,

$$|Pf(re^{i\theta}) - Pf(e^{i\theta_0})| \leq |Pf(re^{i\theta}) - Pf(e^{i\theta})| + |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon.$$

$\Rightarrow Pf$ is continuous at z_0 .



□

$$\left| \int_{-\delta}^{\delta} (f(e^{i(\theta-t)}) - f(e^{i\theta})) P_r(t) dt \right| \leq \int_{-\delta}^{\delta} |f(e^{i(\theta-t)}) - f(e^{i\theta})| P_r(t) dt \leq \int_{-\delta}^{\delta} \frac{\epsilon}{2} P_r(t) dt$$

$$\leq \int_{-\pi}^{\pi} \frac{\epsilon}{2} P_r(t) dt = \frac{\epsilon}{2}.$$

So $1 - \delta' < r < 1 \Rightarrow |Pf(re^{i\theta}) - f(e^{i\theta})| < \epsilon/4 + \epsilon/2 + \epsilon/4 = \epsilon.$
 $\Rightarrow Pf(re^{i\theta}) \rightarrow f(e^{i\theta})$ uniformly in θ as $r \uparrow 1$.

Fix $z_0 = e^{i\theta_0} \in \partial\mathbb{D}$ and $\epsilon > 0$. $f = Pf|_{\partial\mathbb{D}}$ is continuous, so $\exists \delta > 0$ such that

$$|\theta - \theta_0| < \delta \Rightarrow |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon/2$$

For possibly smaller δ


$$1 - \delta < r \leq 1 \Rightarrow |Pf(re^{i\theta}) - Pf(e^{i\theta})| < \epsilon/2 \quad \text{for all } \theta.$$

So if $z = re^{i\theta}$ satisfies $1 - \delta < r \leq 1$ and $|\theta - \theta_0| < \delta$,

$$|Pf(re^{i\theta}) - Pf(e^{i\theta_0})| \leq |Pf(re^{i\theta}) - Pf(e^{i\theta})| + |Pf(e^{i\theta}) - Pf(e^{i\theta_0})| < \epsilon.$$

$\Rightarrow Pf$ is continuous at z_0 .

Remark: We used polar coordinates (homeomorphism near z_0).



□

Translation and scaling gives (proof an exercise):

Corollary

Let $f: \partial\Delta_R(p) \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\Delta_R(p)} \rightarrow \mathbb{R}$, defined by

$$Pf(p + re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(p + Re^{it}) P_{r/R}(\theta - t) dt & \text{if } r < R, \\ f(p + Re^{i\theta}) & \text{if } r = R, \end{cases}$$

is harmonic in $\Delta_R(p)$ and continuous on $\overline{\Delta_R(p)}$.

Translation and scaling gives (proof an exercise):

Corollary

Let $f: \partial\Delta_R(p) \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\Delta_R(p)} \rightarrow \mathbb{R}$, defined by

$$Pf(p + re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(p + Re^{it}) P_{r/R}(\theta - t) dt & \text{if } r < R, \\ f(p + Re^{i\theta}) & \text{if } r = R, \end{cases}$$

is harmonic in $\Delta_R(p)$ and continuous on $\overline{\Delta_R(p)}$.

Represents harmonic functions like Cauchy integral formula:

If f is harmonic on a neighborhood of $\overline{\Delta_R(p)}$ $\Rightarrow P[f|_{\partial\Delta_R(p)}] = f|_{\overline{\Delta_R(p)}}$.

Translation and scaling gives (proof an exercise):

Corollary

Let $f: \partial\Delta_R(p) \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\Delta_R(p)} \rightarrow \mathbb{R}$, defined by

$$Pf(p + re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(p + Re^{it}) P_{r/R}(\theta - t) dt & \text{if } r < R, \\ f(p + Re^{i\theta}) & \text{if } r = R, \end{cases}$$

is harmonic in $\Delta_R(p)$ and continuous on $\overline{\Delta_R(p)}$.

Represents harmonic functions like Cauchy integral formula:

If f is harmonic on a neighborhood of $\overline{\Delta_R(p)}$ $\Rightarrow P[f|_{\partial\Delta_R(p)}] = f|_{\overline{\Delta_R(p)}}$.

However, the Poisson kernel above is specific to the disk.

Translation and scaling gives (proof an exercise):

Corollary

Let $f: \partial\Delta_R(p) \rightarrow \mathbb{R}$ be continuous. Then $Pf: \overline{\Delta_R(p)} \rightarrow \mathbb{R}$, defined by

$$Pf(p + re^{i\theta}) = \begin{cases} \int_{-\pi}^{\pi} f(p + Re^{it}) P_{r/R}(\theta - t) dt & \text{if } r < R, \\ f(p + Re^{i\theta}) & \text{if } r = R, \end{cases}$$

is harmonic in $\Delta_R(p)$ and continuous on $\overline{\Delta_R(p)}$.

Represents harmonic functions like Cauchy integral formula:

If f is harmonic on a neighborhood of $\overline{\Delta_R(p)}$ $\Rightarrow P[f|_{\partial\Delta_R(p)}] = f|_{\overline{\Delta_R(p)}}$.

However, the Poisson kernel above is specific to the disk.

A quick consequence: If f is harmonic on a neighborhood of $\overline{\Delta_R(p)}$, then

$$f(p) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(p + Re^{it}) dt.$$

Poisson kernel works for holomorphic functions (holomorphic functions are harmonic).

Poisson kernel works for holomorphic functions (holomorphic functions are harmonic).

For general functions Cf and Pf are different in the disc:

Consider $f(z) = z + \bar{z} = 2 \operatorname{Re}(z)$ on the unit circle.

$$Pf(z) = z + \bar{z} \quad Cf(z) = z$$

Exercise: Given a bounded continuous $f: \mathbb{R} \rightarrow \mathbb{R}$, prove that $Pf: \overline{\mathbb{H}} \rightarrow \mathbb{R}$,

$$Pf(x + iy) = \begin{cases} \frac{1}{\pi} \int_{-\infty}^{\infty} f(t) \frac{y}{(x-t)^2 + y^2} dt & \text{if } y > 0, \\ f(x) & \text{if } y = 0, \end{cases}$$

is harmonic in \mathbb{H} and continuous on $\overline{\mathbb{H}}$.

Exercise: Given a bounded continuous $f: \mathbb{R} \rightarrow \mathbb{R}$, prove that $Pf: \overline{\mathbb{H}} \rightarrow \mathbb{R}$,

$$Pf(x + iy) = \begin{cases} \frac{1}{\pi} \int_{-\infty}^{\infty} f(t) \frac{y}{(x-t)^2 + y^2} dt & \text{if } y > 0, \\ f(x) & \text{if } y = 0, \end{cases}$$

is harmonic in \mathbb{H} and continuous on $\overline{\mathbb{H}}$.

Exercise: Derive the *Schwarz integral formula*, which recovers a holomorphic function out of the real parts of the boundary values and the value of the imaginary part at one point.

If $f: \overline{\mathbb{D}} \rightarrow \mathbb{C}$ is continuous and holomorphic on \mathbb{D} , then for all $z \in \mathbb{D}$,

$$f(z) = \frac{1}{2\pi i} \int_{\partial \mathbb{D}} \frac{\zeta + z}{\zeta - z} \frac{\operatorname{Re} f(\zeta)}{\zeta} d\zeta + i \operatorname{Im} f(0).$$