

# Cultivating Complex Analysis: Automorphisms of the disc (3.5.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

For  $a \in \mathbb{D}$ , define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

For  $a \in \mathbb{D}$ , define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

### Proposition

For every  $a \in \mathbb{D}$ ,

(i)  $\varphi_a(a) = 0$ ,  $\varphi_a(0) = -a$ ,  $\varphi'_a(0) = 1 - |a|^2$ ,  $\varphi'_a(a) = \frac{1}{1 - |a|^2}$ ,

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

For  $a \in \mathbb{D}$ , define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

### Proposition

For every  $a \in \mathbb{D}$ ,

- (i)  $\varphi_a(a) = 0$ ,  $\varphi_a(0) = -a$ ,  $\varphi'_a(0) = 1 - |a|^2$ ,  $\varphi'_a(a) = \frac{1}{1 - |a|^2}$ ,
- (ii)  $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$ , and  $\varphi_a(\mathbb{D}) = \mathbb{D}$ ,

Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

For  $a \in \mathbb{D}$ , define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$

### Proposition

For every  $a \in \mathbb{D}$ ,

- (i)  $\varphi_a(a) = 0$ ,  $\varphi_a(0) = -a$ ,  $\varphi'_a(0) = 1 - |a|^2$ ,  $\varphi'_a(a) = \frac{1}{1 - |a|^2}$ ,
- (ii)  $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$ , and  $\varphi_a(\mathbb{D}) = \mathbb{D}$ ,
- (iii)  $\varphi_a$  restricted to  $\mathbb{D}$  is an automorphism of the disc and  $\varphi_a^{-1} = \varphi_{-a}$ .



Recall, an automorphism of  $U$  is a biholomorphism  $f: U \rightarrow U$  (bijective such that  $f$  and  $f^{-1}$  are both holomorphic).

Let us compute the automorphism group of the disc,  $\text{Aut}(\mathbb{D})$ .

We start with certain specific automorphisms.

For  $a \in \mathbb{D}$ , define

$$\varphi_a(z) \stackrel{\text{def}}{=} \frac{z - a}{1 - \bar{a}z}.$$


### Proposition

For every  $a \in \mathbb{D}$ ,

- (i)  $\varphi_a(a) = 0$ ,  $\varphi_a(0) = -a$ ,  $\varphi'_a(0) = 1 - |a|^2$ ,  $\varphi'_a(a) = \frac{1}{1 - |a|^2}$ ,
- (ii)  $\varphi_a(\partial\mathbb{D}) = \partial\mathbb{D}$ , and  $\varphi_a(\mathbb{D}) = \mathbb{D}$ ,
- (iii)  $\varphi_a$  restricted to  $\mathbb{D}$  is an automorphism of the disc and  $\varphi_a^{-1} = \varphi_{-a}$ .


Proof is an exercise.

Here is what an automorphism  $\varphi_a$  does ( $a = -0.4$ ).



$a$  and  $\varphi_a(a) = 0$  are marked with dots.


Here is what an automorphism  $\varphi_a$  does ( $a = -0.4$ ).



$a$  and  $\varphi_a(a) = 0$  are marked with dots.

It turns out all automorphisms are  $\varphi_a$  up to a rotation:

Here is what an automorphism  $\varphi_a$  does ( $a = -0.4$ ).



$a$  and  $\varphi_a(a) = 0$  are marked with dots.

It turns out all automorphisms are  $\varphi_a$  up to a rotation:

### Proposition

If  $f \in \text{Aut}(\mathbb{D})$ , then there exists an  $a \in \mathbb{D}$  and  $\theta \in \mathbb{R}$  such that

$$f(z) = e^{i\theta} \frac{z - a}{1 - \bar{a}z} = e^{i\theta} \varphi_a(z).$$

**Proof:** Let  $a = f(0)$ .

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .



**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$   
 $\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$   
 $\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D}$



**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$  attains a maximum in  $\mathbb{D}$

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$  attains a maximum in  $\mathbb{D} \Rightarrow h$  is constant

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$  attains a maximum in  $\mathbb{D} \Rightarrow h$  is constant

$\Rightarrow g(z) = \alpha z$  for some constant  $\alpha$ .

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$  attains a maximum in  $\mathbb{D} \Rightarrow h$  is constant

$\Rightarrow g(z) = \alpha z$  for some constant  $\alpha$ . As  $g \in \text{Aut}(\mathbb{D})$ ,  $|\alpha| = 1$  ( $\alpha = e^{i\theta}$ ).

**Proof:** Let  $a = f(0)$ .

$g = \varphi_a \circ f$  is an automorphism with  $g(0) = 0$  as  $\varphi_a(a) = 0$ .

Find a holomorphic  $h(z)$  such that  $g(z) = zh(z)$  (as in the proof of Schwarz).

Schwarz says that for  $z \in \mathbb{D} \setminus \{0\}$ ,  $|h(z)| = \frac{|g(z)|}{|z|} \leq 1$ .

$h$  can have no zeros:  $h(z) = \frac{g(z)}{z}$  cannot be zero for  $z \neq 0$  as  $g$  is injective,  
and it cannot have a zero at  $z = 0$  as  $h(0) = \lim_{z \rightarrow 0} \frac{g(z)}{z} = g'(0) \neq 0$ .

$g$  is biholomorphic  $\Rightarrow g^{-1}$  is continuous  $\Rightarrow g^{-1}(K)$  compact for compact  $K \subset \mathbb{D}$

$\Rightarrow |g(z)| \rightarrow 1$  as  $|z| \rightarrow 1$  (exercise)  $\Rightarrow |h(z)| \rightarrow 1$  as  $|z| \rightarrow 1$

$\Rightarrow |h(z)|$  attains a minimum in  $\mathbb{D} \Rightarrow \left| \frac{1}{h(z)} \right|$  attains a maximum in  $\mathbb{D} \Rightarrow h$  is constant

$\Rightarrow g(z) = \alpha z$  for some constant  $\alpha$ . As  $g \in \text{Aut}(\mathbb{D})$ ,  $|\alpha| = 1$  ( $\alpha = e^{i\theta}$ ).

Apply  $\varphi_{-a}$  to both sides of  $e^{i\theta}z = \varphi_a \circ f$  to find  $f(z) = \varphi_{-a}(ze^{i\theta}) = e^{i\theta} \varphi_{-ae^{-i\theta}}(z)$ .

□

Some good exercises to try:

**Exercise:** Given distinct  $a, b \in \mathbb{D}$ , show that there exists a unique  $f \in \text{Aut}(\mathbb{D})$  such that  $f(a) = b$  and  $f(b) = a$ .

Some good exercises to try:

**Exercise:** Given distinct  $a, b \in \mathbb{D}$ , show that there exists a unique  $f \in \text{Aut}(\mathbb{D})$  such that  $f(a) = b$  and  $f(b) = a$ .

**Exercise:** The automorphisms of  $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$  are of the form  $\frac{az+b}{cz+d}$  for real numbers  $a, b, c, d$  such that  $ad - bc \neq 0$ .

Some good exercises to try:

**Exercise:** Given distinct  $a, b \in \mathbb{D}$ , show that there exists a unique  $f \in \text{Aut}(\mathbb{D})$  such that  $f(a) = b$  and  $f(b) = a$ .

**Exercise:** The automorphisms of  $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$  are of the form  $\frac{az+b}{cz+d}$  for real numbers  $a, b, c, d$  such that  $ad - bc \neq 0$ .

Another fun exercise, a version of which seems to appear on many exams is the following:

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a domain,  $\overline{\mathbb{D}} \subset U$ , and  $f: U \rightarrow \mathbb{C}$  is holomorphic such that  $f(\partial\mathbb{D}) \subset \partial\mathbb{D}$ . Find a formula for  $f$ .



Some good exercises to try:

**Exercise:** Given distinct  $a, b \in \mathbb{D}$ , show that there exists a unique  $f \in \text{Aut}(\mathbb{D})$  such that  $f(a) = b$  and  $f(b) = a$ .

**Exercise:** The automorphisms of  $\mathbb{H} = \{z \in \mathbb{C} : \text{Im } z > 0\}$  are of the form  $\frac{az+b}{cz+d}$  for real numbers  $a, b, c, d$  such that  $ad - bc \neq 0$ .

Another fun exercise, a version of which seems to appear on many exams is the following:

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a domain,  $\overline{\mathbb{D}} \subset U$ , and  $f: U \rightarrow \mathbb{C}$  is holomorphic such that  $f(\partial\mathbb{D}) \subset \partial\mathbb{D}$ . Find a formula for  $f$ .

Hint: The idea is to show that you can divide by finitely many  $\varphi_a(z)$  for various  $a$  until you get something that has no zeros in  $\mathbb{D}$  and will have to be a constant.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some  $z \neq \zeta$ , then  $f \in \text{Aut}(\mathbb{D})$ .

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some  $z \neq \zeta$ , then  $f \in \text{Aut}(\mathbb{D})$ .

Conversely, if  $f \in \text{Aut}(\mathbb{D})$ , then equality holds in both inequalities for all  $z, \zeta \in \mathbb{D}$ .

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \bar{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some  $z \neq \zeta$ , then  $f \in \text{Aut}(\mathbb{D})$ .

Conversely, if  $f \in \text{Aut}(\mathbb{D})$ , then equality holds in both inequalities for all  $z, \zeta \in \mathbb{D}$ .

Proof is an exercise.

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some  $z \neq \zeta$ , then  $f \in \text{Aut}(\mathbb{D})$ .

Conversely, if  $f \in \text{Aut}(\mathbb{D})$ , then equality holds in both inequalities for all  $z, \zeta \in \mathbb{D}$ .

Proof is an exercise.

The Schwarz–Pick lemma gives a bound on the derivative at all points:

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, nonconstant, and  $f(a) = b$ , then

$$|f'(a)| \leq \frac{1 - |b|^2}{1 - |a|^2}.$$

Using automorphisms one proves an “invariant version” of the Schwarz lemma.

### Lemma (Schwarz–Pick)

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, then

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(\zeta)}f(z)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{\zeta}z} \right| \quad \text{and} \quad \frac{|f'(z)|}{1 - |f(z)|^2} \leq \frac{1}{1 - |z|^2} \quad \text{for all } z, \zeta \in \mathbb{D}.$$

If equality holds in one of the inequalities for some  $z \neq \zeta$ , then  $f \in \text{Aut}(\mathbb{D})$ .

Conversely, if  $f \in \text{Aut}(\mathbb{D})$ , then equality holds in both inequalities for all  $z, \zeta \in \mathbb{D}$ .

Proof is an exercise.

The Schwarz–Pick lemma gives a bound on the derivative at all points:

If  $f: \mathbb{D} \rightarrow \mathbb{D}$  is holomorphic, nonconstant, and  $f(a) = b$ , then

$$|f'(a)| \leq \frac{1 - |b|^2}{1 - |a|^2}.$$

Equality  $\Rightarrow f(z) = \varphi_{-b}(e^{i\theta} \varphi_a(z))$  for some  $\theta \in \mathbb{R}$ .