

Cultivating Complex Analysis: Cauchy–Riemann equations (2.1.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

$$Df|_{z_0} = \begin{bmatrix} \frac{\partial u}{\partial x}|_{z_0} & \frac{\partial u}{\partial y}|_{z_0} \\ \frac{\partial v}{\partial x}|_{z_0} & \frac{\partial v}{\partial y}|_{z_0} \end{bmatrix}, \quad \text{where } f = u + iv \quad \text{and} \quad z = x + iy.$$

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

$$Df|_{z_0} = \begin{bmatrix} \frac{\partial u}{\partial x}|_{z_0} & \frac{\partial u}{\partial y}|_{z_0} \\ \frac{\partial v}{\partial x}|_{z_0} & \frac{\partial v}{\partial y}|_{z_0} \end{bmatrix}, \quad \text{where } f = u + iv \quad \text{and} \quad z = x + iy.$$

Question: When does $(Df|_{z_0})h$ correspond to ξh for some $\xi \in \mathbb{C}$?

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

$$Df|_{z_0} = \begin{bmatrix} \frac{\partial u}{\partial x}|_{z_0} & \frac{\partial u}{\partial y}|_{z_0} \\ \frac{\partial v}{\partial x}|_{z_0} & \frac{\partial v}{\partial y}|_{z_0} \end{bmatrix}, \quad \text{where } f = u + iv \quad \text{and} \quad z = x + iy.$$

Question: When does $(Df|_{z_0})h$ correspond to ξh for some $\xi \in \mathbb{C}$?

Answer: When $Df|_{z_0}$ is of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$.

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

$$Df|_{z_0} = \begin{bmatrix} \frac{\partial u}{\partial x}|_{z_0} & \frac{\partial u}{\partial y}|_{z_0} \\ \frac{\partial v}{\partial x}|_{z_0} & \frac{\partial v}{\partial y}|_{z_0} \end{bmatrix}, \quad \text{where } f = u + iv \quad \text{and} \quad z = x + iy.$$

Question: When does $(Df|_{z_0})h$ correspond to ξh for some $\xi \in \mathbb{C}$?

Answer: When $Df|_{z_0}$ is of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$.

I.e., when

$$\frac{\partial u}{\partial x}|_{z_0} = \frac{\partial v}{\partial y}|_{z_0}, \quad \frac{\partial v}{\partial x}|_{z_0} = -\frac{\partial u}{\partial y}|_{z_0}.$$

Let $U \subset \mathbb{C}$ be open. Think of \mathbb{C} as \mathbb{R}^2 for a moment.

Recall that $f: U \rightarrow \mathbb{C}$ is (real) differentiable at $z_0 \in U$ if there exists a real-linear (2×2 matrix) $Df|_{z_0}$ such that

$$\lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - (Df|_{z_0})h|}{|h|} = 0 \quad (\text{again } h \in \mathbb{R}^2 = \mathbb{C}).$$

$$Df|_{z_0} = \begin{bmatrix} \frac{\partial u}{\partial x}|_{z_0} & \frac{\partial u}{\partial y}|_{z_0} \\ \frac{\partial v}{\partial x}|_{z_0} & \frac{\partial v}{\partial y}|_{z_0} \end{bmatrix}, \quad \text{where } f = u + iv \quad \text{and} \quad z = x + iy.$$

Question: When does $(Df|_{z_0})h$ correspond to ξh for some $\xi \in \mathbb{C}$?

Answer: When $Df|_{z_0}$ is of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$.

I.e., when

$$\frac{\partial u}{\partial x}|_{z_0} = \frac{\partial v}{\partial y}|_{z_0}, \quad \frac{\partial v}{\partial x}|_{z_0} = -\frac{\partial u}{\partial y}|_{z_0}.$$

Then $Df|_{z_0}$ corresponds to multiplication by $\xi = \frac{\partial u}{\partial x}|_{z_0} + i\frac{\partial v}{\partial x}|_{z_0} = \frac{\partial v}{\partial y}|_{z_0} - i\frac{\partial u}{\partial y}|_{z_0}$.

Consequently,

$$0 = \lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - \xi h|}{|h|} = \lim_{h \rightarrow 0} \left| \frac{f(z_0 + h) - f(z_0)}{h} - \xi \right|,$$

Consequently,

$$0 = \lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - \xi h|}{|h|} = \lim_{h \rightarrow 0} \left| \frac{f(z_0 + h) - f(z_0)}{h} - \xi \right|,$$

or

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \xi.$$

Consequently,

$$0 = \lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - \xi h|}{|h|} = \lim_{h \rightarrow 0} \left| \frac{f(z_0 + h) - f(z_0)}{h} - \xi \right|,$$

or

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \xi.$$

So f is complex differentiable at z_0 and $f'(z_0) = \xi$.

Consequently,

$$0 = \lim_{h \rightarrow 0} \frac{|f(z_0 + h) - f(z_0) - \xi h|}{|h|} = \lim_{h \rightarrow 0} \left| \frac{f(z_0 + h) - f(z_0)}{h} - \xi \right|,$$

or

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \xi.$$

So f is complex differentiable at z_0 and $f'(z_0) = \xi$.

We proved that: If f is (real) differentiable at z_0 , with $\frac{\partial u}{\partial x}\big|_{z_0} = \frac{\partial v}{\partial y}\big|_{z_0}$ and $\frac{\partial v}{\partial x}\big|_{z_0} = -\frac{\partial u}{\partial y}\big|_{z_0}$, then f is complex differentiable at z_0 .

Conversely, if f is complex differentiable at z_0 , then it is real differentiable at z_0 (the complex derivative $f'(z_0)$ gives the $Df|_{z_0}$).

Conversely, if f is complex differentiable at z_0 , then it is real differentiable at z_0 (the complex derivative $f'(z_0)$ gives the $Df|_{z_0}$).

Further, $\frac{\partial u}{\partial x}\big|_{z_0} = \frac{\partial v}{\partial y}\big|_{z_0}$ and $\frac{\partial v}{\partial x}\big|_{z_0} = -\frac{\partial u}{\partial y}\big|_{z_0}$ must also hold.

Conversely, if f is complex differentiable at z_0 , then it is real differentiable at z_0 (the complex derivative $f'(z_0)$ gives the $Df|_{z_0}$).

Further, $\frac{\partial u}{\partial x}|_{z_0} = \frac{\partial v}{\partial y}|_{z_0}$ and $\frac{\partial v}{\partial x}|_{z_0} = -\frac{\partial u}{\partial y}|_{z_0}$ must also hold.

Proposition

Let $U \subset \mathbb{C}$ be open and $f = u + iv: U \rightarrow \mathbb{C}$ be a function. Then f is complex differentiable at $z_0 \in U$ if and only if f (real) differentiable at $z_0 \in U$ with $\frac{\partial u}{\partial x}|_{z_0} = \frac{\partial v}{\partial y}|_{z_0}$ and $\frac{\partial v}{\partial x}|_{z_0} = -\frac{\partial u}{\partial y}|_{z_0}$.

Conversely, if f is complex differentiable at z_0 , then it is real differentiable at z_0 (the complex derivative $f'(z_0)$ gives the $Df|_{z_0}$).

Further, $\frac{\partial u}{\partial x}\big|_{z_0} = \frac{\partial v}{\partial y}\big|_{z_0}$ and $\frac{\partial v}{\partial x}\big|_{z_0} = -\frac{\partial u}{\partial y}\big|_{z_0}$ must also hold.

Proposition

Let $U \subset \mathbb{C}$ be open and $f = u + iv: U \rightarrow \mathbb{C}$ be a function. Then f is complex differentiable at $z_0 \in U$ if and only if f (real) differentiable at $z_0 \in U$ with $\frac{\partial u}{\partial x}\big|_{z_0} = \frac{\partial v}{\partial y}\big|_{z_0}$ and $\frac{\partial v}{\partial x}\big|_{z_0} = -\frac{\partial u}{\partial y}\big|_{z_0}$.

In this case, $f'(z_0) = \frac{\partial u}{\partial x}\big|_{z_0} + i\frac{\partial v}{\partial x}\big|_{z_0} = \frac{\partial v}{\partial y}\big|_{z_0} - i\frac{\partial u}{\partial y}\big|_{z_0}$.

If the partial derivatives exist and are continuous, then f is real differentiable, so:

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable).

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable). Then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad (1)$$

if and only if f is holomorphic (complex differentiable at all $z \in U$),

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable). Then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad (1)$$

if and only if f is holomorphic (complex differentiable at all $z \in U$), or in other words,

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \quad \text{exists for all } z \in U.$$

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable). Then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad (1)$$

if and only if f is holomorphic (complex differentiable at all $z \in U$), or in other words,

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \quad \text{exists for all } z \in U.$$

The equations (1) are called the *Cauchy–Riemann equations*.

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable). Then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad (1)$$

if and only if f is holomorphic (complex differentiable at all $z \in U$), or in other words,

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \quad \text{exists for all } z \in U.$$

The equations (1) are called the *Cauchy–Riemann equations*.

Complex analysis is the study of their solutions.

If the partial derivatives exist and are continuous, then f is real differentiable, so:

Corollary

Let $U \subset \mathbb{C}$ be open and let $f = u + iv: U \rightarrow \mathbb{C}$ be a function such that $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, and $\frac{\partial v}{\partial y}$ exist and are continuous (that is, f is continuously differentiable). Then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad (1)$$

if and only if f is holomorphic (complex differentiable at all $z \in U$), or in other words,

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \quad \text{exists for all } z \in U.$$

The equations (1) are called the *Cauchy–Riemann equations*.

Complex analysis is the study of their solutions.

Remark: If only the partial derivatives exist but aren't continuous, the function may fail to be differentiable (or even continuous) and may not be holomorphic even if it satisfies (1).

Exercise: Show that e^z is holomorphic and its complex derivative is e^z , and hence $\sin z$ and $\cos z$ are also holomorphic.

Exercise: Show that e^z is holomorphic and its complex derivative is e^z , and hence $\sin z$ and $\cos z$ are also holomorphic.

Hint: Note that $e^{x+iy} = e^x \cos y + ie^x \sin y$ and use the Corollary from previous slide.