

Cultivating Complex Analysis:

Types of singularities and Riemann extension (5.2.1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

A holomorphic $f: U \setminus \{p\} \rightarrow \mathbb{C}$ has an *isolated singularity* at p .

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

A holomorphic $f: U \setminus \{p\} \rightarrow \mathbb{C}$ has an *isolated singularity* at p .

An isolated singularity is *removable* if there exists a holomorphic $F: U \rightarrow \mathbb{C}$ such that $f(z) = F(z)$ for all $z \in U \setminus \{p\}$.

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

A holomorphic $f: U \setminus \{p\} \rightarrow \mathbb{C}$ has an *isolated singularity* at p .

An isolated singularity is *removable* if there exists a holomorphic $F: U \rightarrow \mathbb{C}$ such that $f(z) = F(z)$ for all $z \in U \setminus \{p\}$.

An isolated singularity p is a *pole* if $\lim_{z \rightarrow p} f(z) = \infty$.

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

A holomorphic $f: U \setminus \{p\} \rightarrow \mathbb{C}$ has an *isolated singularity* at p .

An isolated singularity is *removable* if there exists a holomorphic $F: U \rightarrow \mathbb{C}$ such that $f(z) = F(z)$ for all $z \in U \setminus \{p\}$.

An isolated singularity p is a *pole* if $\lim_{z \rightarrow p} f(z) = \infty$.

An isolated singularity that is neither removable nor a pole is an *essential singularity*.

Definition

Suppose $U \subset \mathbb{C}$ is open and $p \in U$.

A holomorphic $f: U \setminus \{p\} \rightarrow \mathbb{C}$ has an *isolated singularity* at p .

An isolated singularity is *removable* if there exists a holomorphic $F: U \rightarrow \mathbb{C}$ such that $f(z) = F(z)$ for all $z \in U \setminus \{p\}$.

An isolated singularity p is a *pole* if $\lim_{z \rightarrow p} f(z) = \infty$.

An isolated singularity that is neither removable nor a pole is an *essential singularity*.

Examples: Pole: $1/z$, essential: $e^{1/z}$.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z)$$

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z) = 0.$$

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z) = 0.$$

So g is complex differentiable at p and so holomorphic on U .

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z) = 0.$$

So g is complex differentiable at p and so holomorphic on U .

As $g(p) = 0$ and $g'(p) = 0$, g has a zero of order $k \geq 2$.

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z) = 0.$$

So g is complex differentiable at p and so holomorphic on U .

As $g(p) = 0$ and $g'(p) = 0$, g has a zero of order $k \geq 2$.

Write $g(z) = (z - p)^k h(z)$, where h is holomorphic on U .

Theorem (Riemann extension theorem)

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ is holomorphic. If f is bounded (near p suffices), then p is a removable singularity.

Proof: Let $g(z) = (z - p)^2 f(z)$ for $z \neq p$ and $g(p) = 0$.

g is holomorphic in $U \setminus \{p\}$.

Supposing f is bounded,

$$\lim_{z \rightarrow p} \frac{g(z) - g(p)}{z - p} = \lim_{z \rightarrow p} (z - p)f(z) = 0.$$

So g is complex differentiable at p and so holomorphic on U .

As $g(p) = 0$ and $g'(p) = 0$, g has a zero of order $k \geq 2$.

Write $g(z) = (z - p)^k h(z)$, where h is holomorphic on U .

Then $f(z) = (z - p)^{k-2} h(z)$, that is, p is a removable singularity.



Exercise: Prove that if $f: \mathbb{D} \setminus \{0\} \rightarrow \mathbb{D} \setminus \{0\}$ is an automorphism, then $f(z) = e^{i\theta}z$ for some θ .

Exercise: Prove that if $f: \mathbb{D} \setminus \{0\} \rightarrow \mathbb{D} \setminus \{0\}$ is an automorphism, then $f(z) = e^{i\theta}z$ for some θ .

The Riemann extension theorem is (of course) not true for functions that are not holomorphic.

Exercise: Prove that if $f: \mathbb{D} \setminus \{0\} \rightarrow \mathbb{D} \setminus \{0\}$ is an automorphism, then $f(z) = e^{i\theta}z$ for some θ .

The Riemann extension theorem is (of course) not true for functions that are not holomorphic.

Exercise: Prove that $\frac{xy}{x^2+y^2}$ is a bounded infinitely (real) differentiable function on $\mathbb{R}^2 \setminus \{(0,0)\}$ with an isolated singularity, and this function does not extend through the singularity even continuously.

Poles are places where the function “blows up” to a finite order:

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .*

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Then f is not zero in some $\Delta_r(p) \setminus \{p\}$ (it goes to infinity).

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Then f is not zero in some $\Delta_r(p) \setminus \{p\}$ (it goes to infinity).

$1/f$ is bounded in some $\Delta_r(p) \setminus \{p\}$ (as $f \rightarrow \infty$).

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Then f is not zero in some $\Delta_r(p) \setminus \{p\}$ (it goes to infinity).

$1/f$ is bounded in some $\Delta_r(p) \setminus \{p\}$ (as $f \rightarrow \infty$).

Thus, by Riemann extension, there exists an h holomorphic in $\Delta_r(p)$ such that $h(z) = 1/f(z)$ for $z \neq p$.

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Then f is not zero in some $\Delta_r(p) \setminus \{p\}$ (it goes to infinity).

$1/f$ is bounded in some $\Delta_r(p) \setminus \{p\}$ (as $f \rightarrow \infty$).

Thus, by Riemann extension, there exists an h holomorphic in $\Delta_r(p)$ such that $h(z) = 1/f(z)$ for $z \neq p$.

As $\frac{1}{f} \rightarrow 0$ as $z \rightarrow p$, we have $h(p) = 0$.

Poles are places where the function “blows up” to a finite order:

Corollary

Suppose $U \subset \mathbb{C}$ is open, $p \in U$, and $f: U \setminus \{p\} \rightarrow \mathbb{C}$ holomorphic.

- (i) If p is a pole, then there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ has a removable singularity at p .
- (ii) Conversely, if there exists a $k \in \mathbb{N}$ such that $g(z) = (z - p)^k f(z)$ is bounded near p (has a removable singularity), then f has a pole or a removable singularity at p .

Proof: Suppose f has a pole at p .

Then f is not zero in some $\Delta_r(p) \setminus \{p\}$ (it goes to infinity).

$1/f$ is bounded in some $\Delta_r(p) \setminus \{p\}$ (as $f \rightarrow \infty$).

Thus, by Riemann extension, there exists an h holomorphic in $\Delta_r(p)$ such that $h(z) = 1/f(z)$ for $z \neq p$.

As $\frac{1}{f} \rightarrow 0$ as $z \rightarrow p$, we have $h(p) = 0$.

$\Rightarrow h(z) = (z - p)^k \psi(z)$ for some holomorphic ψ and $k \in \mathbb{N}$, where $\psi(p) \neq 0$.

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z - p)^k \psi(z)$ $g(z) = (z - p)^k f(z)$)

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p) \Rightarrow 1/\psi$ is holomorphic in $\Delta_r(p)$.

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p) \Rightarrow 1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z)$$

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p) \Rightarrow 1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)}$$

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p)$ \Rightarrow $1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p) \Rightarrow 1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

For the converse statement suppose $g(z)$ has a removable singularity.

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p)$ \Rightarrow $1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

For the converse statement suppose $g(z)$ has a removable singularity.

Then $g(z) = (z-p)^\ell \varphi(z)$ where $\varphi(p) \neq 0$.

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p)$ \Rightarrow $1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

For the converse statement suppose $g(z)$ has a removable singularity.

Then $g(z) = (z-p)^\ell \varphi(z)$ where $\varphi(p) \neq 0$.

Then

$$f(z) = (z-p)^{\ell-k} \varphi(z),$$

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p)$ \Rightarrow $1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

For the converse statement suppose $g(z)$ has a removable singularity.

Then $g(z) = (z-p)^\ell \varphi(z)$ where $\varphi(p) \neq 0$.

Then

$$f(z) = (z-p)^{\ell-k} \varphi(z),$$

$f(z)$ either goes to ∞ if $k > \ell$ (f has a pole) or

(keeping track: $h(z) = 1/f(z)$ $h(z) = (z-p)^k \psi(z)$ $g(z) = (z-p)^k f(z)$)

As $h \neq 0$ in $\Delta_r(p) \setminus \{p\}$, then $\psi \neq 0$ in $\Delta_r(p) \Rightarrow 1/\psi$ is holomorphic in $\Delta_r(p)$.

For $z \in \Delta_r(p) \setminus \{p\}$,

$$g(z) = (z-p)^k f(z) = (z-p)^k \frac{1}{(z-p)^k \psi(z)} = \frac{1}{\psi(z)}.$$

and (i) is done.

For the converse statement suppose $g(z)$ has a removable singularity.

Then $g(z) = (z-p)^\ell \varphi(z)$ where $\varphi(p) \neq 0$.

Then

$$f(z) = (z-p)^{\ell-k} \varphi(z),$$

$f(z)$ either goes to ∞ if $k > \ell$ (f has a pole) or
is bounded near p if $k \leq \ell$ (f has a removable singularity).



Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole.

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Symmetry between zeros and poles:

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Symmetry between zeros and poles:

If f has a zero of order k at p , then $1/f$ has a pole of order k at p .

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Symmetry between zeros and poles:

If f has a zero of order k at p , then $1/f$ has a pole of order k at p .

If f has a pole of order k at p , then $1/f$ has a removable singularity, and the extended function has a zero of order k at p .

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Symmetry between zeros and poles:

If f has a zero of order k at p , then $1/f$ has a pole of order k at p .

If f has a pole of order k at p , then $1/f$ has a removable singularity, and the extended function has a zero of order k at p .

More precisely, if f has a pole or a removable singularity we can write $f(z) = (z - p)^\ell g(z)$ for some $\ell \in \mathbb{Z}$ and some holomorphic g such that $g(p) \neq 0$.

Definition

Given a holomorphic function f with a pole at p , the smallest $k \in \mathbb{N}$ such that $(z - p)^k f(z)$ is bounded near p is called the *order* of the pole. A pole of order 1 is called a *simple pole*.

So if f has a pole at p , then

$$f(z) = \frac{g(z)}{(z - p)^k}$$

for holomorphic g nonzero at p , and k is the order of the pole.

Symmetry between zeros and poles:

If f has a zero of order k at p , then $1/f$ has a pole of order k at p .

If f has a pole of order k at p , then $1/f$ has a removable singularity, and the extended function has a zero of order k at p .

More precisely, if f has a pole or a removable singularity we can write $f(z) = (z - p)^\ell g(z)$ for some $\ell \in \mathbb{Z}$ and some holomorphic g such that $g(p) \neq 0$.

p is a zero of order ℓ if $\ell > 0$, and p is a pole of order $-\ell$ if $\ell < 0$.

A useful exercise that parallels a result we proved for zeros:

A useful exercise that parallels a result we proved for zeros:

Exercise: Suppose f has a pole of order $k \in \mathbb{N}$ at p . Show that there exists a holomorphic g defined near p such that $g(p) = 0$ and $g'(p) \neq 0$ and such that near p

$$f(z) = \frac{1}{(g(z))^k}.$$