

Cultivating Complex Analysis: The Riemann sphere (1.3)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

Real numbers are sometimes extended to add $\pm\infty$.

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

To define topology, define a bijection $g: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ by

$$g(z) = \begin{cases} 1/z & \text{if } z \neq 0 \text{ and } z \neq \infty, \\ \infty & \text{if } z = 0, \\ 0 & \text{if } z = \infty. \end{cases}$$

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

To define topology, define a bijection $g: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ by

$$g(z) = \begin{cases} 1/z & \text{if } z \neq 0 \text{ and } z \neq \infty, \\ \infty & \text{if } z = 0, \\ 0 & \text{if } z = \infty. \end{cases}$$

We want all open sets in \mathbb{C} to be open in \mathbb{C}_∞ .

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

To define topology, define a bijection $g: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ by

$$g(z) = \begin{cases} 1/z & \text{if } z \neq 0 \text{ and } z \neq \infty, \\ \infty & \text{if } z = 0, \\ 0 & \text{if } z = \infty. \end{cases}$$

We want all open sets in \mathbb{C} to be open in \mathbb{C}_∞ .

Neighborhoods of ∞ are $g(U)$ for neighborhoods U of 0.

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

To define topology, define a bijection $g: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ by

$$g(z) = \begin{cases} 1/z & \text{if } z \neq 0 \text{ and } z \neq \infty, \\ \infty & \text{if } z = 0, \\ 0 & \text{if } z = \infty. \end{cases}$$

We want all open sets in \mathbb{C} to be open in \mathbb{C}_∞ .

Neighborhoods of ∞ are $g(U)$ for neighborhoods U of 0.

If talking about convergence at $z \in \mathbb{C}$, we just use the normal topology.

Real numbers are sometimes extended to add $\pm\infty$.

Similarly, the complex plane is extended by adding ∞ (only one infinity):

$$\mathbb{C}_\infty = \mathbb{C} \cup \{\infty\}.$$

We call \mathbb{C}_∞ the *Riemann sphere*.

To define topology, define a bijection $g: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$ by

$$g(z) = \begin{cases} 1/z & \text{if } z \neq 0 \text{ and } z \neq \infty, \\ \infty & \text{if } z = 0, \\ 0 & \text{if } z = \infty. \end{cases}$$

We want all open sets in \mathbb{C} to be open in \mathbb{C}_∞ .

Neighborhoods of ∞ are $g(U)$ for neighborhoods U of 0.

If talking about convergence at $z \in \mathbb{C}$, we just use the normal topology.

If talking about convergence at ∞ , we map with g and consider the topology at 0.

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

Start with $\Phi((0, 0, 1)) = \infty$.

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

Start with $\Phi((0, 0, 1)) = \infty$.

For any $p \in S^2 \setminus (0, 0, 1)$ there is a unique line through $(0, 0, 1)$ and p .

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

Start with $\Phi((0, 0, 1)) = \infty$.

For any $p \in S^2 \setminus (0, 0, 1)$ there is a unique line through $(0, 0, 1)$ and p .

The line intersects the xy -plane at a unique point $(x, y, 0)$.

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:

Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

Start with $\Phi((0, 0, 1)) = \infty$.

For any $p \in S^2 \setminus (0, 0, 1)$ there is a unique line through $(0, 0, 1)$ and p .

The line intersects the xy -plane at a unique point $(x, y, 0)$.

We identify $(x, y, 0)$ with $\xi = x + iy$ and define $\Phi(p) = \xi$.

We can give \mathbb{C}_∞ a metric space structure through the *stereographic projection*:


Let S^2 be the unit sphere in \mathbb{R}^3 . We define a bijection $\Phi: S^2 \rightarrow \mathbb{C}_\infty$.

Start with $\Phi((0,0,1)) = \infty$.

For any $p \in S^2 \setminus (0,0,1)$ there is a unique line through $(0,0,1)$ and p .

The line intersects the xy -plane at a unique point $(x,y,0)$.

We identify $(x,y,0)$ with $\xi = x + iy$ and define $\Phi(p) = \xi$.



The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

$$L \in \mathbb{C} \text{ and } z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}.$$

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

$$L \in \mathbb{C} \text{ and } z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}.$$

$$L \in \mathbb{C}: \quad \lim_{z \rightarrow \infty} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall \epsilon > 0 \exists M \text{ such that } |f(z) - L| < \epsilon \text{ whenever } |z| > M.$$

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

$$L \in \mathbb{C} \text{ and } z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}.$$

$$L \in \mathbb{C}: \quad \lim_{z \rightarrow \infty} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall \epsilon > 0 \exists M \text{ such that } |f(z) - L| < \epsilon \text{ whenever } |z| > M.$$

$$z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = \infty \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall M > 0 \exists \delta > 0 \text{ such that } |f(z)| > M \text{ whenever } |z - z_0| < \delta.$$

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

$$L \in \mathbb{C} \text{ and } z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}.$$

$$L \in \mathbb{C}: \quad \lim_{z \rightarrow \infty} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall \epsilon > 0 \exists M \text{ such that } |f(z) - L| < \epsilon \text{ whenever } |z| > M.$$

$$z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = \infty \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall M > 0 \exists \delta > 0 \text{ such that } |f(z)| > M \text{ whenever } |z - z_0| < \delta.$$

$$L \in \mathbb{C}_\infty: \quad \lim_{z \rightarrow \infty} f(z) = L \Leftrightarrow \lim_{z \rightarrow 0} f(1/z) = L.$$

The point of the Riemann sphere is to give the value ∞ to certain limits and to allow limits as z tends to ∞ .

Note that conveniently \mathbb{C}_∞ is compact (one-point compactification of \mathbb{C}).

For $f: U \subset \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$, define

$$\lim_{z \rightarrow z_0} f(z) = L$$

using the topology of the Riemann sphere.

It is an exercise to show the following:

$$L \in \mathbb{C} \text{ and } z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \lim_{z \rightarrow z_0} f(z) = L \text{ in } \mathbb{C}.$$

$$L \in \mathbb{C}: \quad \lim_{z \rightarrow \infty} f(z) = L \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall \epsilon > 0 \exists M \text{ such that } |f(z) - L| < \epsilon \text{ whenever } |z| > M.$$

$$z_0 \in \mathbb{C}: \quad \lim_{z \rightarrow z_0} f(z) = \infty \text{ in } \mathbb{C}_\infty \Leftrightarrow \forall M > 0 \exists \delta > 0 \text{ such that } |f(z)| > M \text{ whenever } |z - z_0| < \delta.$$

$$L \in \mathbb{C}_\infty: \quad \lim_{z \rightarrow \infty} f(z) = L \Leftrightarrow \lim_{z \rightarrow 0} f(1/z) = L.$$

$$z_0 \in \mathbb{C}_\infty: \quad \lim_{z \rightarrow z_0} f(z) = \infty \Leftrightarrow \lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

To see it, prove that for large z , $|P(z)| \geq c |z|^d$ for some c .

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

To see it, prove that for large z , $|P(z)| \geq c |z|^d$ for some c .

So every nonconstant polynomial is ∞ at ∞ .

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

To see it, prove that for large z , $|P(z)| \geq c |z|^d$ for some c .

So every nonconstant polynomial is ∞ at ∞ .

Remark: Be careful about limits in extended real sense vs. Riemann sphere sense:

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does not exist,} \quad \text{but} \quad \lim_{z \rightarrow 0} \frac{1}{z} = \infty.$$

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

To see it, prove that for large z , $|P(z)| \geq c |z|^d$ for some c .

So every nonconstant polynomial is ∞ at ∞ .

Remark: Be careful about limits in extended real sense vs. Riemann sphere sense:

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does not exist,} \quad \text{but} \quad \lim_{z \rightarrow 0} \frac{1}{z} = \infty.$$

(if we think of x =real, and z =complex.)

It makes sense to talk about the value of $1/z$ at the origin as ∞ , and the value of z at ∞ as ∞ .

Consider a polynomial $P(z) = a_d z^d + a_{d-1} z^{d-1} + \cdots + a_1 z + a_0$.

If $d \geq 1$ and $a_d \neq 0$, then it is an exercise to show that

$$\lim_{z \rightarrow \infty} P(z) = \infty$$

To see it, prove that for large z , $|P(z)| \geq c |z|^d$ for some c .

So every nonconstant polynomial is ∞ at ∞ .

Remark: Be careful about limits in extended real sense vs. Riemann sphere sense:

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does not exist,} \quad \text{but} \quad \lim_{z \rightarrow 0} \frac{1}{z} = \infty.$$

(if we think of x =real, and z =complex.)

We'll use Riemann sphere sense unless otherwise noted or obvious. We may use $+\infty$ to distinguish from Riemann sphere ∞ if confusion could arise.

Partial arithmetic can be defined on \mathbb{C}_∞ :

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Neither $\infty + \infty$ nor $\infty - \infty$ makes sense:

if $f(z) = z$ and $g(z) = -z$, then $\lim_{z \rightarrow \infty} f(z) = \infty$, $\lim_{z \rightarrow \infty} g(z) = \infty$, but $\lim_{z \rightarrow \infty} (f(z) + g(z)) = 0$.

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Neither $\infty + \infty$ nor $\infty - \infty$ makes sense:

if $f(z) = z$ and $g(z) = -z$, then $\lim_{z \rightarrow \infty} f(z) = \infty$, $\lim_{z \rightarrow \infty} g(z) = \infty$, but $\lim_{z \rightarrow \infty} (f(z) + g(z)) = 0$.

If $c \neq 0$, then define $c/0 = \infty$ and $c \cdot \infty = \infty$.

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Neither $\infty + \infty$ nor $\infty - \infty$ makes sense:

if $f(z) = z$ and $g(z) = -z$, then $\lim_{z \rightarrow \infty} f(z) = \infty$, $\lim_{z \rightarrow \infty} g(z) = \infty$, but $\lim_{z \rightarrow \infty} (f(z) + g(z)) = 0$.

If $c \neq 0$, then define $c/0 = \infty$ and $c \cdot \infty = \infty$.

If $c \neq \infty$, then define $c/\infty = 0$.

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Neither $\infty + \infty$ nor $\infty - \infty$ makes sense:

if $f(z) = z$ and $g(z) = -z$, then $\lim_{z \rightarrow \infty} f(z) = \infty$, $\lim_{z \rightarrow \infty} g(z) = \infty$, but $\lim_{z \rightarrow \infty} (f(z) + g(z)) = 0$.

If $c \neq 0$, then define $c/0 = \infty$ and $c \cdot \infty = \infty$.

If $c \neq \infty$, then define $c/\infty = 0$.

$0 \cdot \infty$, $0/0$, and ∞/∞ are undefined.

Partial arithmetic can be defined on \mathbb{C}_∞ :

If $c \neq \infty$, declare $c + \infty = \infty$.

Neither $\infty + \infty$ nor $\infty - \infty$ makes sense:

if $f(z) = z$ and $g(z) = -z$, then $\lim_{z \rightarrow \infty} f(z) = \infty$, $\lim_{z \rightarrow \infty} g(z) = \infty$, but $\lim_{z \rightarrow \infty} (f(z) + g(z)) = 0$.

If $c \neq 0$, then define $c/0 = \infty$ and $c \cdot \infty = \infty$.

If $c \neq \infty$, then define $c/\infty = 0$.

$0 \cdot \infty$, $0/0$, and ∞/∞ are undefined.

Note that this is different from the extended real arithmetic.