

Cultivating Complex Analysis: Matrix representation of complex numbers (1.1.4)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

\mathbb{C} is a 2-dimensional real vector space.

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

1 " = " $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and i " = " $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ are identity and rotation counterclockwise by 90° .

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

1 " = " $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and i " = " $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ are identity and rotation counterclockwise by 90° .

\mathbb{C} can be identified as a subring of $M_2(\mathbb{R})$:

Multiplying the matrices is multiplying the numbers (exercise).

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

1 " = " $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and i " = " $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ are identity and rotation counterclockwise by 90° .

\mathbb{C} can be identified as a subring of $M_2(\mathbb{R})$:

Multiplying the matrices is multiplying the numbers (exercise).

Complex conjugation is real-linear, $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, but not a multiplication by a complex number.

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

1 " " $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and i " " $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ are identity and rotation counterclockwise by 90° .

\mathbb{C} can be identified as a subring of $M_2(\mathbb{R})$:

Multiplying the matrices is multiplying the numbers (exercise).

Complex conjugation is real-linear, $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, but not a multiplication by a complex number.

Multiplication by complex numbers and complex conjugation "generate" the ring $M_2(\mathbb{R})$:

All real-linear operators are of the form $z \mapsto \xi z + \zeta \bar{z}$ (exercise).

\mathbb{C} is a 2-dimensional real vector space.

Multiplication $z \mapsto \xi z$ is a real-linear operator $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ if $\xi = a + ib$:

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} ac - bd \\ bc + ad \end{bmatrix} \quad \text{just like} \quad (a + ib)(c + id) = (ac - bd) + i(bc + ad).$$

1 " " $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and i " " $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ are identity and rotation counterclockwise by 90° .

\mathbb{C} can be identified as a subring of $M_2(\mathbb{R})$:

Multiplying the matrices is multiplying the numbers (exercise).

Complex conjugation is real-linear, $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, but not a multiplication by a complex number.

Multiplication by complex numbers and complex conjugation "generate" the ring $M_2(\mathbb{R})$:

All real-linear operators are of the form $z \mapsto \xi z + \zeta \bar{z}$ (exercise).

Easy exercise: If M is $z \mapsto \xi z$, then $\det(M) = |\xi|^2$. If M is $z \mapsto \xi z + \zeta \bar{z}$, then $\det(M) = |\xi|^2 - |\zeta|^2$.

How will we use this representation of \mathbb{C} ?

How will we use this representation of \mathbb{C} ?

A differentiable $f: \mathbb{C} \rightarrow \mathbb{C}$ is really $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

How will we use this representation of \mathbb{C} ?

A differentiable $f: \mathbb{C} \rightarrow \mathbb{C}$ is really $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

The derivative of f is a linear operator from \mathbb{R}^2 to \mathbb{R}^2 .

How will we use this representation of \mathbb{C} ?

A differentiable $f: \mathbb{C} \rightarrow \mathbb{C}$ is really $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

The derivative of f is a linear operator from \mathbb{R}^2 to \mathbb{R}^2 .

Complex analysis is the study of functions (holomorphic) whose derivative is multiplication by a complex number.

How will we use this representation of \mathbb{C} ?

A differentiable $f: \mathbb{C} \rightarrow \mathbb{C}$ is really $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

The derivative of f is a linear operator from \mathbb{R}^2 to \mathbb{R}^2 .

Complex analysis is the study of functions (holomorphic) whose derivative is multiplication by a complex number.

The derivative is a matrix of partial derivatives, so to be of the form $\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ is a system of partial differential equations (PDE): the Cauchy–Riemann equations.