

# Cultivating Complex Analysis: Hurwitz's theorem (5.4.3)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

**Proof:**  $\Gamma$  is compact  $\Rightarrow$  there is a  $\delta > 0$  such that  $\delta < |f(z)|$  for all  $z \in \Gamma$ .

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

**Proof:**  $\Gamma$  is compact  $\Rightarrow$  there is a  $\delta > 0$  such that  $\delta < |f(z)|$  for all  $z \in \Gamma$ .

$\{f_n\}$  converges uniformly to  $f$  on  $\Gamma$ .

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

**Proof:**  $\Gamma$  is compact  $\Rightarrow$  there is a  $\delta > 0$  such that  $\delta < |f(z)|$  for all  $z \in \Gamma$ .

$\{f_n\}$  converges uniformly to  $f$  on  $\Gamma$ .

For  $n$  large enough,

$$|f(z) - f_n(z)| < \delta < |f(z)| \quad \forall z \in \Gamma$$

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

**Proof:**  $\Gamma$  is compact  $\Rightarrow$  there is a  $\delta > 0$  such that  $\delta < |f(z)|$  for all  $z \in \Gamma$ .

$\{f_n\}$  converges uniformly to  $f$  on  $\Gamma$ .

For  $n$  large enough,

$$|f(z) - f_n(z)| < \delta < |f(z)| \quad \forall z \in \Gamma$$

Rouché's theorem  $\Rightarrow$   $f$  and  $f_n$  have the same number of zeros in  $V$ . □

## Theorem (Hurwitz)

Let  $U \subset \mathbb{C}$  be open and  $f_n: U \rightarrow \mathbb{C}$  a sequence of holomorphic functions converging uniformly on compact subsets to a holomorphic  $f: U \rightarrow \mathbb{C}$ . Suppose  $\Gamma$  is a cycle in  $U$  homologous to zero in  $U$ , such that  $n(\Gamma; z)$  is 0 or 1 for all  $z \notin \Gamma$ . Suppose  $f$  has no zeros on  $\Gamma$  and  $k$  zeros (counting multiplicity) in  $V = \{z \in U \setminus \Gamma : n(\Gamma; z) = 1\}$ . Then there is an  $N$  such that for all  $n \geq N$ ,  $f_n$  has  $k$  zeros (counting multiplicity) in  $V$ .

**Proof:**  $\Gamma$  is compact  $\Rightarrow$  there is a  $\delta > 0$  such that  $\delta < |f(z)|$  for all  $z \in \Gamma$ .

$\{f_n\}$  converges uniformly to  $f$  on  $\Gamma$ .

For  $n$  large enough,

$$|f(z) - f_n(z)| < \delta < |f(z)| \quad \forall z \in \Gamma$$

Rouché's theorem  $\Rightarrow$   $f$  and  $f_n$  have the same number of zeros in  $V$ . □

“No zeros on  $\Gamma$ ” is necessary:  $\Gamma = \partial\mathbb{D}$ ,  $f(z) = z - 1$ ,  $f_n(z) = z + (1 - \frac{1}{n})$ .

**Example:** For every integer  $k > 0$ ,  $\exists N$  such that  $\forall d \geq N$ ,

$$P_d(z) = \sum_{n=0}^d \frac{(-1)^n}{(2n)!} z^{2n}$$

has exactly  $2k$  zeros in  $\Delta_{\pi k}(0)$ .



**Example:** For every integer  $k > 0$ ,  $\exists N$  such that  $\forall d \geq N$ ,

$$P_d(z) = \sum_{n=0}^d \frac{(-1)^n}{(2n)!} z^{2n}$$

has exactly  $2k$  zeros in  $\Delta_{\pi k}(0)$ .

**Proof:**

$P_d$  are the partial sums of the power series of  $\cos(z)$ , which has exactly  $2k$  zeros in  $\Delta_{\pi k}(0)$ .

The usual application is for a small disc:

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ .

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

Then for a small enough  $\Delta_r(z_0)$ ,  $\exists N$  such that  $\forall n \geq N$ ,  
 $f_n$  has  $k$  zeros counting multiplicity in  $\Delta_r(z_0)$ .  
(use  $\Gamma = \partial\Delta_r(z_0)$ )

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

Then for a small enough  $\Delta_r(z_0)$ ,  $\exists N$  such that  $\forall n \geq N$ ,  
 $f_n$  has  $k$  zeros counting multiplicity in  $\Delta_r(z_0)$ .  
(use  $\Gamma = \partial\Delta_r(z_0)$ )

Hurwitz theorem does not work for real functions.

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

Then for a small enough  $\Delta_r(z_0)$ ,  $\exists N$  such that  $\forall n \geq N$ ,  
 $f_n$  has  $k$  zeros counting multiplicity in  $\Delta_r(z_0)$ .  
(use  $\Gamma = \partial\Delta_r(z_0)$ )

Hurwitz theorem does not work for real functions.

**Example:**

$f(x) = x^2$ ,  $f_n(x) = x^2 + \frac{1}{n}$ .  $f_n \rightarrow f$  uniformly, but  $f_n$  never zero.

The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

Then for a small enough  $\Delta_r(z_0)$ ,  $\exists N$  such that  $\forall n \geq N$ ,

$f_n$  has  $k$  zeros counting multiplicity in  $\Delta_r(z_0)$ .

(use  $\Gamma = \partial\Delta_r(z_0)$ )

Hurwitz theorem does not work for real functions.

**Example:**

$f(x) = x^2$ ,  $f_n(x) = x^2 + \frac{1}{n}$ .  $f_n \rightarrow f$  uniformly, but  $f_n$  never zero.

$f(z) = z^2$ ,  $f_n(z) = z^2 + \frac{1}{n}$ .  $f_n \rightarrow f$  uniformly.



The usual application is for a small disc:

Suppose  $\{f_n\}$  is a sequence of holomorphic functions on  $U$  converging uniformly on compact subsets to  $f$ . Suppose  $z_0$  is a zero of  $f$  of order  $k$ .

Then for a small enough  $\Delta_r(z_0)$ ,  $\exists N$  such that  $\forall n \geq N$ ,  
 $f_n$  has  $k$  zeros counting multiplicity in  $\Delta_r(z_0)$ .  
(use  $\Gamma = \partial\Delta_r(z_0)$ )

Hurwitz theorem does not work for real functions.

**Example:**

$f(x) = x^2$ ,  $f_n(x) = x^2 + \frac{1}{n}$ .  $f_n \rightarrow f$  uniformly, but  $f_n$  never zero.

$f(z) = z^2$ ,  $f_n(z) = z^2 + \frac{1}{n}$ .  $f_n \rightarrow f$  uniformly.

For any  $\epsilon > 0$ ,  $z^2 + \frac{1}{n}$  has two zeros in  $\Delta_\epsilon(0)$ , for large enough  $n$ :  $\pm i\sqrt{\frac{1}{n}}$ .

Injective holomorphic mappings are called *univalent*.

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant.

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,



Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,

such that  $f - w$  is not zero on  $\overline{\Delta_r(z_1)} \setminus \{z_1\}$  or  $\overline{\Delta_r(z_2)} \setminus \{z_2\}$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,

such that  $f - w$  is not zero on  $\overline{\Delta_r(z_1)} \setminus \{z_1\}$  or  $\overline{\Delta_r(z_2)} \setminus \{z_2\}$ .

Hurwitz  $\Rightarrow$  for large enough  $n$ ,  $f_n - w$  has the same number of zeros in  $\Delta_r(z_1)$  as  $f - w$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

*Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.*

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,

such that  $f - w$  is not zero on  $\overline{\Delta_r(z_1)} \setminus \{z_1\}$  or  $\overline{\Delta_r(z_2)} \setminus \{z_2\}$ .

Hurwitz  $\Rightarrow$  for large enough  $n$ ,  $f_n - w$  has the same number of zeros in  $\Delta_r(z_1)$  as  $f - w$ .  
Same for  $\Delta_r(z_2)$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,

such that  $f - w$  is not zero on  $\overline{\Delta_r(z_1)} \setminus \{z_1\}$  or  $\overline{\Delta_r(z_2)} \setminus \{z_2\}$ .

Hurwitz  $\Rightarrow$  for large enough  $n$ ,  $f_n - w$  has the same number of zeros in  $\Delta_r(z_1)$  as  $f - w$ .  
Same for  $\Delta_r(z_2)$ .

$\Rightarrow \exists z'_1 \in \Delta_r(z_1)$  and  $z'_2 \in \Delta_r(z_2)$  such that  $f_n(z'_1) = f_n(z'_2) = w$ .

Injective holomorphic mappings are called *univalent*.

### Corollary

Suppose  $U \subset \mathbb{C}$  is a domain and  $f_n: U \rightarrow \mathbb{C}$  are univalent holomorphic functions that converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Then  $f$  is either univalent or constant.

**Proof:** Suppose  $f$  is nonconstant. Suppose  $\exists$  distinct  $z_1$  and  $z_2$  in  $U$  such that  $f(z_1) = f(z_2) = w$ .

$f - w$  has isolated zeros at  $z_1$  and  $z_2$ .

Take  $\overline{\Delta_r(z_1)}, \overline{\Delta_r(z_2)} \subset U$ ,

$\Delta_r(z_1) \cap \Delta_r(z_2) = \emptyset$ ,

such that  $f - w$  is not zero on  $\overline{\Delta_r(z_1)} \setminus \{z_1\}$  or  $\overline{\Delta_r(z_2)} \setminus \{z_2\}$ .

Hurwitz  $\Rightarrow$  for large enough  $n$ ,  $f_n - w$  has the same number of zeros in  $\Delta_r(z_1)$  as  $f - w$ .  
Same for  $\Delta_r(z_2)$ .

$\Rightarrow \exists z'_1 \in \Delta_r(z_1)$  and  $z'_2 \in \Delta_r(z_2)$  such that  $f_n(z'_1) = f_n(z'_2) = w$ .

$\Rightarrow f_n$  not univalent.



Another common application is the following exercise.

Another common application is the following exercise.

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a domain, and  $f_n: U \rightarrow \mathbb{C}$  are holomorphic, nowhere zero, and converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ .

Another common application is the following exercise.

**Exercise:** Suppose  $U \subset \mathbb{C}$  is a domain, and  $f_n: U \rightarrow \mathbb{C}$  are holomorphic, nowhere zero, and converge uniformly on compact subsets to  $f: U \rightarrow \mathbb{C}$ . Show that either  $f$  is nowhere zero, or  $f$  is identically zero.