

# Cultivating Complex Analysis: Montel's theorem (6.2)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

## Definition

Let  $U \subset \mathbb{C}$  be open. A set  $\mathcal{F}$  of holomorphic functions  $f: U \rightarrow \mathbb{C}$  is called a *normal family* if every sequence in  $\mathcal{F}$  has a subsequence that converges uniformly on compact subsets (the limit need not be in  $\mathcal{F}$ ).

## Definition

Let  $U \subset \mathbb{C}$  be open. A set  $\mathcal{F}$  of holomorphic functions  $f: U \rightarrow \mathbb{C}$  is called a *normal family* if every sequence in  $\mathcal{F}$  has a subsequence that converges uniformly on compact subsets (the limit need not be in  $\mathcal{F}$ ).

A set  $\mathcal{F}$  of functions on  $U$  is *locally bounded* if for every  $p \in U$ , there is a disc  $\Delta_r(p) \subset U$  and  $M > 0$  such that  $\|f\|_{\Delta_r(p)} \leq M$  for all  $f \in \mathcal{F}$ .

## Definition

Let  $U \subset \mathbb{C}$  be open. A set  $\mathcal{F}$  of holomorphic functions  $f: U \rightarrow \mathbb{C}$  is called a *normal family* if every sequence in  $\mathcal{F}$  has a subsequence that converges uniformly on compact subsets (the limit need not be in  $\mathcal{F}$ ).

A set  $\mathcal{F}$  of functions on  $U$  is *locally bounded* if for every  $p \in U$ , there is a disc  $\Delta_r(p) \subset U$  and  $M > 0$  such that  $\|f\|_{\Delta_r(p)} \leq M$  for all  $f \in \mathcal{F}$  (i.e.,  $|f(z)| \leq M$  for all  $z \in \Delta_r(p)$  and all  $f \in \mathcal{F}$ ).

## Definition

Let  $U \subset \mathbb{C}$  be open. A set  $\mathcal{F}$  of holomorphic functions  $f: U \rightarrow \mathbb{C}$  is called a *normal family* if every sequence in  $\mathcal{F}$  has a subsequence that converges uniformly on compact subsets (the limit need not be in  $\mathcal{F}$ ).

A set  $\mathcal{F}$  of functions on  $U$  is *locally bounded* if for every  $p \in U$ , there is a disc  $\Delta_r(p) \subset U$  and  $M > 0$  such that  $\|f\|_{\Delta_r(p)} \leq M$  for all  $f \in \mathcal{F}$  (i.e.,  $|f(z)| \leq M$  for all  $z \in \Delta_r(p)$  and all  $f \in \mathcal{F}$ ).

In more modern language:

$\mathcal{F}$  normal family

=

$\mathcal{F}$  precompact in the space of holomorphic functions on  $U$  with the topology of uniform convergence on compact subsets.

## Definition

Let  $U \subset \mathbb{C}$  be open. A set  $\mathcal{F}$  of holomorphic functions  $f: U \rightarrow \mathbb{C}$  is called a *normal family* if every sequence in  $\mathcal{F}$  has a subsequence that converges uniformly on compact subsets (the limit need not be in  $\mathcal{F}$ ).

A set  $\mathcal{F}$  of functions on  $U$  is *locally bounded* if for every  $p \in U$ , there is a disc  $\Delta_r(p) \subset U$  and  $M > 0$  such that  $\|f\|_{\Delta_r(p)} \leq M$  for all  $f \in \mathcal{F}$  (i.e.,  $|f(z)| \leq M$  for all  $z \in \Delta_r(p)$  and all  $f \in \mathcal{F}$ ).

In more modern language:

$\mathcal{F}$  normal family

=

$\mathcal{F}$  precompact in the space of holomorphic functions on  $U$  with the topology of uniform convergence on compact subsets.

**Exercise:** Prove that “locally bounded” means “bounded on compact subsets.”

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family*

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*



## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ .

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right|$$

## Theorem (Montel)

*Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).*

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta|$$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta|$$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right|$$



## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta|$$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$  is equicontinuous ( $\frac{4M}{r}$  does not depend on  $f$ ).

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$  is equicontinuous ( $\frac{4M}{r}$  does not depend on  $f$ ).

$\Rightarrow$  Arzelà–Ascoli applies to any sequence in  $\mathcal{F}$

## Theorem (Montel)

Let  $U \subset \mathbb{C}$  be open and let  $\mathcal{F}$  be a locally bounded set of holomorphic functions on  $U$ . Then  $\mathcal{F}$  is a normal family (every sequence has a subsequence that converges uniformly on compact subsets).

We apply Arzelà–Ascoli, so we prove  $\mathcal{F}$  being holomorphic means  $\mathcal{F}$  is equicontinuous.

**Proof:** Consider  $p \in U$ ,  $\overline{\Delta_r(p)} \subset U$ , such that  $\|f\|_{\overline{\Delta_r(p)}} \leq M$  for all  $f \in \mathcal{F}$ .

If  $z \in \overline{\Delta_{r/2}(p)}$  and  $\zeta \in \partial\Delta_r(p)$ , then  $|\zeta - z| \geq r/2$ . So for  $z \in \overline{\Delta_{r/2}(p)}$  and all  $f \in \mathcal{F}$ ,

$$|f'(z)| = \left| \frac{1}{2\pi i} \int_{\partial\Delta_r(p)} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta \right| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{|f(\zeta)|}{|\zeta - z|^2} |d\zeta| \leq \frac{1}{2\pi} \int_{\partial\Delta_r(p)} \frac{M}{(r/2)^2} |d\zeta| = \frac{4M}{r}.$$

$$|f(z) - f(p)| = \left| \int_{[p,z]} f'(\zeta) d\zeta \right| \leq \int_{[p,z]} |f'(\zeta)| |d\zeta| \leq \frac{4M}{r} |z - p|.$$

$\Rightarrow \mathcal{F}$  is equicontinuous ( $\frac{4M}{r}$  does not depend on  $f$ ).

$\Rightarrow$  Arzelà–Ascoli applies to any sequence in  $\mathcal{F} \Rightarrow \mathcal{F}$  is a normal family. □

Montel's theorem is good for finding extremal functions.

Montel's theorem is good for finding extremal functions.  
E.g., we will prove the Riemann mapping theorem:

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected  $U$  to  $\mathbb{D}$  is a map that maximizes the derivative at a point.



Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected  $U$  to  $\mathbb{D}$  is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected  $U$  to  $\mathbb{D}$  is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Another commonly used consequence of Montel is Vitali's theorem.

### Theorem (Vitali)

*Suppose  $U \subset \mathbb{C}$  is a domain,  $\{f_n\}$  is a locally bounded sequence of holomorphic functions that converges pointwise on a set  $E \subset U$ , and  $E$  has a limit point in  $U$ . Then  $\{f_n\}$  converges uniformly on compact subsets in  $U$ .*

Montel's theorem is good for finding extremal functions.

E.g., we will prove the Riemann mapping theorem: A biholomorphism of a simply connected  $U$  to  $\mathbb{D}$  is a map that maximizes the derivative at a point.

Montel gives us a way of finding a maximizer.

Another commonly used consequence of Montel is Vitali's theorem.

### Theorem (Vitali)

*Suppose  $U \subset \mathbb{C}$  is a domain,  $\{f_n\}$  is a locally bounded sequence of holomorphic functions that converges pointwise on a set  $E \subset U$ , and  $E$  has a limit point in  $U$ . Then  $\{f_n\}$  converges uniformly on compact subsets in  $U$ .*

Proof is an exercise.

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family.

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family. Note: The converse is false without an extra hypothesis.

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family. Note: The converse is false without an extra hypothesis.

**Exercise:** Let  $U \subset \mathbb{C}$  be a domain,  $p \in U$ , and suppose  $\exists$  a nonconstant bounded holomorphic function on  $U$ .

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family. Note: The converse is false without an extra hypothesis.

**Exercise:** Let  $U \subset \mathbb{C}$  be a domain,  $p \in U$ , and suppose  $\exists$  a nonconstant bounded holomorphic function on  $U$ .

a) Prove  $\exists$  a holomorphic  $F: U \rightarrow \mathbb{D}$  such that  $F'(p) \neq 0$ , and  $|f'(p)| \leq |F'(p)|$  for all holomorphic  $f: U \rightarrow \mathbb{D}$ .



**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family. Note: The converse is false without an extra hypothesis.

**Exercise:** Let  $U \subset \mathbb{C}$  be a domain,  $p \in U$ , and suppose  $\exists$  a nonconstant bounded holomorphic function on  $U$ .

- a) Prove  $\exists$  a holomorphic  $F: U \rightarrow \mathbb{D}$  such that  $F'(p) \neq 0$ , and  $|f'(p)| \leq |F'(p)|$  for all holomorphic  $f: U \rightarrow \mathbb{D}$ .
- b) Show  $F(p) = 0$ .

**Exercise:** Prove the converse to Montel: If  $\mathcal{F}$  is a normal family of holomorphic functions on an open set  $U \subset \mathbb{C}$ , then  $\mathcal{F}$  is locally bounded.

**Exercise:** Let  $U \subset \mathbb{C}$  be open and  $\mathcal{F}$  a normal family of holomorphic functions on  $U$ . Show that  $\{f' : f \in \mathcal{F}\}$  is a normal family. Note: The converse is false without an extra hypothesis.

**Exercise:** Let  $U \subset \mathbb{C}$  be a domain,  $p \in U$ , and suppose  $\exists$  a nonconstant bounded holomorphic function on  $U$ .

- a) Prove  $\exists$  a holomorphic  $F: U \rightarrow \mathbb{D}$  such that  $F'(p) \neq 0$ , and  $|f'(p)| \leq |F'(p)|$  for all holomorphic  $f: U \rightarrow \mathbb{D}$ .
- b) Show  $F(p) = 0$ .

**Exercise:** Show that if the partial sums of a power series centered at  $p$  are uniformly bounded on  $\Delta_r(p)$  for some  $r > 0$ , then the power series converges in  $\Delta_r(p)$ .