

# Cultivating Complex Analysis: Linear fractional transformations (1.4 part 1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on  $a, b, c, d$  guarantees that the ratio does not simplify.

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on  $a, b, c, d$  guarantees that the ratio does not simplify.

If  $c = 0$ ,  $f$  can be written as  $f(z) = az + b$ .

Define  $f(\infty) = \infty$ .

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on  $a, b, c, d$  guarantees that the ratio does not simplify.

If  $c = 0$ ,  $f$  can be written as  $f(z) = az + b$ .

Define  $f(\infty) = \infty$ .

If  $c \neq 0$ , the expression is defined on  $\mathbb{C} \setminus \{-d/c\}$ .

Define  $f(-d/c) = \infty$  and  $f(\infty) = \frac{a}{c}$ .

$$f(z) = \frac{az + b}{cz + d} \quad \text{where } ad \neq bc$$

is a *linear fractional transformation* (LFT), (or *Möbius transformation*).

The condition on  $a, b, c, d$  guarantees that the ratio does not simplify.

If  $c = 0$ ,  $f$  can be written as  $f(z) = az + b$ .

Define  $f(\infty) = \infty$ .

If  $c \neq 0$ , the expression is defined on  $\mathbb{C} \setminus \{-d/c\}$ .

Define  $f(-d/c) = \infty$  and  $f(\infty) = \frac{a}{c}$ .

So takes the Riemann sphere to Riemann sphere.

It is an easy exercise that  $f: \mathbb{C}_\infty \rightarrow \mathbb{C}_\infty$  is bijective and continuous.

Any LFT is a composition of:

Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$



Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

*complex dilations*

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

*complex dilations*

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

*complex dilations*

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Consider an LFT  $f(z) = \frac{az+b}{cz+d}$ . WLOG  $c = 1$  or  $c = 0$ .

Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

*complex dilations*

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

and *inversions*

$$I(z) = \frac{1}{z}.$$

Consider an LFT  $f(z) = \frac{az+b}{cz+d}$ . WLOG  $c = 1$  or  $c = 0$ .

If  $c = 1$ , then

$$f(z) = \frac{az + b}{z + d} = \frac{b - ad}{z + d} + a = T_a \left( D_{b-ad} \left( I(T_d(z)) \right) \right).$$

Any LFT is a composition of:

*translations*

$$T_a(z) = z + a, \quad a \in \mathbb{C},$$

*complex dilations*

$$D_a(z) = az, \quad a \in \mathbb{C} \setminus \{0\},$$

*and inversions*

$$I(z) = \frac{1}{z}.$$

Consider an LFT  $f(z) = \frac{az+b}{cz+d}$ . WLOG  $c = 1$  or  $c = 0$ .

If  $c = 1$ , then

$$f(z) = \frac{az + b}{z + d} = \frac{b - ad}{z + d} + a = T_a \left( D_{b-ad} \left( I(T_d(z)) \right) \right).$$

If  $c = 0$ , assume  $d = 1$  and  $f(z) = az + b$ :

$$f(z) = az + b = T_b(D_a(z)).$$

A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .

A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .

Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .

Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.



A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .


Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.



A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .

Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.


The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.

A use for this decomposition (exercise):

*LFTs take the set of straight lines and circles to the set of straight lines and circles.*



A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .

Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:


$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.

A use for this decomposition (exercise):

*LFTs take the set of straight lines and circles to the set of straight lines and circles.*

In fact, any straight line or circle can be taken to any other straight line or circle.



A translation  $T_a$  just moves everything in  $\mathbb{C}$  in the  $a$  direction and fixes  $\infty$ .


Complex dilation  $D_a$  is the normal euclidean dilation (scaling) by  $|a|$  and rotation by  $\arg a$ . It also fixes  $\infty$ .

The inversion is the euclidean plane geometry inversion across the unit circle and then a conjugation.

The euclidean inversion inverts the distance:

$$\frac{1}{|z|} e^{i \arg z} = \frac{|z|}{|z|^2} e^{i \arg z} = \frac{1}{\bar{z}}.$$

To get complex inversion we conjugate.



A use for this decomposition (exercise):

*LFTs take the set of straight lines and circles to the set of straight lines and circles.*

In fact, any straight line or circle can be taken to any other straight line or circle.

**Remark:** A straight line is really a very large circle through  $\infty$ .