

# Cultivating Complex Analysis: Cauchy's formula in a disc (3.2.4)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

### Theorem (Cauchy integral formula in a disc)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $\overline{\Delta_r(p)} \subset U$ . Then for all  $z \in \Delta_r(p)$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

### Theorem (Cauchy integral formula in a disc)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $\overline{\Delta_r(p)} \subset U$ . Then for all  $z \in \Delta_r(p)$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

### Theorem (Cauchy integral formula in a disc)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $\overline{\Delta_r(p)} \subset U$ . Then for all  $z \in \Delta_r(p)$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

### Theorem (Cauchy integral formula in a disc)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $\overline{\Delta_r(p)} \subset U$ . Then for all  $z \in \Delta_r(p)$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)

are given in terms of values on the boundary (small set).

We get to perhaps the most fundamental theorem in complex analysis:  
The Cauchy integral formula.

We prove a version in a disc.

### Theorem (Cauchy integral formula in a disc)

*Suppose  $U \subset \mathbb{C}$  is open,  $f: U \rightarrow \mathbb{C}$  is holomorphic,  $\overline{\Delta_r(p)} \subset U$ . Then for all  $z \in \Delta_r(p)$ ,*

$$f(z) = \frac{1}{2\pi i} \int_{\partial \Delta_r(p)} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

It should be surprising:

Values inside (large set)

are given in terms of values on the boundary (small set).

A quick (but hardly only) application is to compute integrals of expressions such as  $\frac{\cos(z^2)}{z(z-1)}$  that blow up somewhere inside the cycle.



**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ .


**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ . Write  $\alpha = \partial\Delta_s(z)$ .

**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ . Write  $\alpha = \partial\Delta_s(z)$ .

Connect  $\alpha$  to  $\gamma$  via two straight lines  
to give two closed paths  $c_1$  and  $c_2$




**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ . Write  $\alpha = \partial\Delta_s(z)$ .

Connect  $\alpha$  to  $\gamma$  via two straight lines to give two closed paths  $c_1$  and  $c_2$

As chains,  $c_1 + c_2 = \gamma - \alpha$ .



**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.


Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ . Write  $\alpha = \partial\Delta_s(z)$ .

Connect  $\alpha$  to  $\gamma$  via two straight lines to give two closed paths  $c_1$  and  $c_2$

As chains,  $c_1 + c_2 = \gamma - \alpha$ .

Each  $c_j$  lies in a star-like domain

where  $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$  is holomorphic.



**Proof:** Fix  $z \in \Delta_r(p)$ , and write  $\gamma = \partial\Delta_r(p)$  oriented counterclockwise.

Let  $\Delta_s(z)$  be a small disc where  $\overline{\Delta_s(z)} \subset \Delta_r(p)$ . Write  $\alpha = \partial\Delta_s(z)$ .

Connect  $\alpha$  to  $\gamma$  via two straight lines to give two closed paths  $c_1$  and  $c_2$

As chains,  $c_1 + c_2 = \gamma - \alpha$ .


Each  $c_j$  lies in a star-like domain

where  $\zeta \mapsto \frac{f(\zeta)}{\zeta - z}$  is holomorphic. So

$$\int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta =$$

$$\int_{c_1} \frac{f(\zeta)}{\zeta - z} d\zeta + \int_{c_2} \frac{f(\zeta)}{\zeta - z} d\zeta = 0 + 0.$$

(by Cauchy's theorem)



So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$



So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough,

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough, so we can take the limit as  $s \rightarrow 0$ .

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough, so we can take the limit as  $s \rightarrow 0$ .

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough, so we can take the limit as  $s \rightarrow 0$ .

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta$$



So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough, so we can take the limit as  $s \rightarrow 0$ .

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt$$

So

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

Parametrize  $\alpha$  as  $\alpha(t) = z + se^{it}$ .

$$\frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(z + se^{it})}{z + se^{it} - z} sie^{it} dt = \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt.$$

The integral over  $\gamma$  (independent of  $s$ ) equals the integral over  $\alpha$  for all  $s > 0$  small enough, so we can take the limit as  $s \rightarrow 0$ .

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi i} \int_{\alpha} \frac{f(\zeta)}{\zeta - z} d\zeta = \lim_{s \downarrow 0} \frac{1}{2\pi} \int_0^{2\pi} f(z + se^{it}) dt = f(z).$$

(by continuity of  $f$  at  $z$ ) □

There is a very useful consequence left as an easy exercise:

There is a very useful consequence left as an easy exercise:

**Exercise:** Suppose  $f$  is holomorphic in an open neighborhood of  $\overline{\Delta_r(p)}$ . Show that  $f$  at  $p$  is the average of the values on  $\partial\Delta_r(p)$ . That is, show

$$f(p) = \frac{1}{2\pi} \int_0^{2\pi} f(p + re^{it}) dt.$$