

Cultivating Complex Analysis: Line integrals (3.1 part 1)

Jiří Lebl

Departemento pri Matematiko de Oklahoma Ŝtata Universitato

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A path γ is *simple closed* if $\gamma(a) = \gamma(b)$ and $\gamma|_{(a,b]}$ is injective.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A path γ is *simple closed* if $\gamma(a) = \gamma(b)$ and $\gamma|_{(a,b]}$ is injective.

Piecewise- C^1 means that \exists numbers $t_0 = a < t_1 < \cdots < t_k = b$ such that $\gamma|_{[t_{\ell-1}, t_\ell]}$ is C^1 (up to the endpoints) and its derivative is never zero.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A path γ is *simple closed* if $\gamma(a) = \gamma(b)$ and $\gamma|_{(a,b]}$ is injective.

Piecewise- C^1 means that \exists numbers $t_0 = a < t_1 < \cdots < t_k = b$ such that $\gamma|_{[t_{\ell-1}, t_\ell]}$ is C^1 (up to the endpoints) and its derivative is never zero.

Equivalently, γ is C^1 on every $(t_{\ell-1}, t_\ell)$, γ' is never zero, and

$$\lim_{t \uparrow t_\ell} \gamma'(t) \quad \lim_{t \downarrow t_\ell} \gamma'(t)$$

exist and are never zero.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A path γ is *simple closed* if $\gamma(a) = \gamma(b)$ and $\gamma|_{(a,b]}$ is injective.

Piecewise- C^1 means that \exists numbers $t_0 = a < t_1 < \cdots < t_k = b$ such that $\gamma|_{[t_{\ell-1}, t_\ell]}$ is C^1 (up to the endpoints) and its derivative is never zero.

Equivalently, γ is C^1 on every $(t_{\ell-1}, t_\ell)$, γ' is never zero, and

$$\lim_{t \uparrow t_\ell} \gamma'(t) \quad \lim_{t \downarrow t_\ell} \gamma'(t)$$

exist and are never zero.

We also consider γ as a set, we write γ to mean $\gamma([a, b])$.

A *piecewise- C^1 path* or a *path* for short is a continuous complex-valued piecewise continuously differentiable function $\gamma: [a, b] \rightarrow \mathbb{C}$ such that $\gamma'(t)$ and all its one-sided limits are never 0.

A path γ is *closed* if $\gamma(a) = \gamma(b)$.

A path γ is *simple closed* if $\gamma(a) = \gamma(b)$ and $\gamma|_{(a,b]}$ is injective.

Piecewise- C^1 means that \exists numbers $t_0 = a < t_1 < \cdots < t_k = b$ such that $\gamma|_{[t_{\ell-1}, t_\ell]}$ is C^1 (up to the endpoints) and its derivative is never zero.

Equivalently, γ is C^1 on every $(t_{\ell-1}, t_\ell)$, γ' is never zero, and

$$\lim_{t \uparrow t_\ell} \gamma'(t) \quad \lim_{t \downarrow t_\ell} \gamma'(t)$$


exist and are never zero.

We also consider γ as a set, we write γ to mean $\gamma([a, b])$.

E.g., we say γ is in U or $\gamma \subset U$ if $\gamma([a, b]) \subset U$.


Example: Consider $\gamma: [0, 4] \rightarrow \mathbb{C}$,

$$\gamma(t) = \begin{cases} t & \text{if } t \in [0, 1], \\ 1 + i(t - 1) & \text{if } t \in (1, 2], \\ 3 - t + i & \text{if } t \in (2, 3], \\ i(4 - t) & \text{if } t \in (3, 4]. \end{cases}$$



Example: Consider $\gamma: [0, 4] \rightarrow \mathbb{C}$,


$$\gamma(t) = \begin{cases} t & \text{if } t \in [0, 1], \\ 1 + i(t - 1) & \text{if } t \in (1, 2], \\ 3 - t + i & \text{if } t \in (2, 3], \\ i(4 - t) & \text{if } t \in (3, 4]. \end{cases}$$



Note, for example, on $t \in (0, 1)$, $\gamma'(t) = 1$, and so $\lim_{t \uparrow 1} \gamma'(t) = 1$.

Example: Consider $\gamma: [0, 4] \rightarrow \mathbb{C}$,

$$\gamma(t) = \begin{cases} t & \text{if } t \in [0, 1], \\ 1 + i(t - 1) & \text{if } t \in (1, 2], \\ 3 - t + i & \text{if } t \in (2, 3], \\ i(4 - t) & \text{if } t \in (3, 4]. \end{cases}$$



Note, for example, on $t \in (0, 1)$, $\gamma'(t) = 1$, and so $\lim_{t \uparrow 1} \gamma'(t) = 1$.

Similarly $\lim_{t \downarrow 1} \gamma'(t) = i$, etc.

Given a piecewise- C^1 path $\gamma: [a, b] \rightarrow \mathbb{C}$ and a continuous function f on γ , we define the *line integral* (or *path integral*, *curve integral*, *contour integral*)

$$\int_{\gamma} f(z) dz \stackrel{\text{def}}{=} \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

Given a piecewise- C^1 path $\gamma: [a, b] \rightarrow \mathbb{C}$ and a continuous function f on γ , we define the *line integral* (or *path integral*, *curve integral*, *contour integral*)

$$\int_{\gamma} f(z) dz \stackrel{\text{def}}{=} \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

The RHS makes sense: The integrand is bounded and continuous except at finitely many points, so Riemann integrable.


Given a piecewise- C^1 path $\gamma: [a, b] \rightarrow \mathbb{C}$ and a continuous function f on γ , we define the *line integral* (or *path integral*, *curve integral*, *contour integral*)

$$\int_{\gamma} f(z) dz \stackrel{\text{def}}{=} \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

The RHS makes sense: The integrand is bounded and continuous except at finitely many points, so Riemann integrable.

The definition makes sense even if $\gamma'(t)$ is zero somewhere.


Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.



Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$




Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.




Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz$$




Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt$$




Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt = ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt.$$



Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.


For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt = ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt.$$

When $n + 1 = 0$, the integral is 2π and $r^{n+1} = 1$.



Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim


$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$

First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt = ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt.$$

When $n + 1 = 0$, the integral is 2π and $r^{n+1} = 1$.

Other n are a calculus exercise.



Example: Let $\gamma: [0, 2\pi] \rightarrow \mathbb{C}$ given by $\gamma(t) = re^{it}$ be the circle of radius r oriented counterclockwise: $\partial\Delta_r(0)$.

For $n \in \mathbb{Z}$, we claim

$$\int_{\gamma} z^n dz = \begin{cases} 2\pi i & \text{if } n = -1, \\ 0 & \text{otherwise.} \end{cases}$$


First, $\gamma'(t) = ire^{it}$.

$$\int_{\gamma} z^n dz = \int_0^{2\pi} r^n e^{int} ire^{it} dt = ir^{n+1} \int_0^{2\pi} e^{i(n+1)t} dt.$$

When $n + 1 = 0$, the integral is 2π and $r^{n+1} = 1$.

Other n are a calculus exercise.

Note that the value of the integral does not depend on r .



The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

Just write $dz = dx + i dy$, which means writing $\gamma(t) = x(t) + i y(t)$:

$$\int_{\gamma} f(z) dz$$

The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

Just write $dz = dx + i dy$, which means writing $\gamma(t) = x(t) + i y(t)$:

$$\int_{\gamma} f(z) dz = \int_{\gamma} f(z) (dx + i dy)$$

The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

Just write $dz = dx + i dy$, which means writing $\gamma(t) = x(t) + i y(t)$:

$$\int_{\gamma} f(z) dz = \int_{\gamma} f(z) (dx + i dy) = \int_{\gamma} f(z) dx + i f(z) dy$$

The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

Just write $dz = dx + i dy$, which means writing $\gamma(t) = x(t) + i y(t)$:

$$\int_{\gamma} f(z) dz = \int_{\gamma} f(z) (dx + i dy) = \int_{\gamma} f(z) dx + i \int_{\gamma} f(z) dy = \int_a^b \underbrace{\left(f(\gamma(t))x'(t) + i f(\gamma(t))y'(t) \right)}_{f(\gamma(t))\gamma'(t)} dt.$$

The definition is the same as the one you've seen in multivariable calculus,

$$\int_{\gamma} P dx + Q dy.$$

Just write $dz = dx + i dy$, which means writing $\gamma(t) = x(t) + i y(t)$:

$$\int_{\gamma} f(z) dz = \int_{\gamma} f(z) (dx + i dy) = \int_{\gamma} f(z) dx + i f(z) dy = \int_a^b \underbrace{\left(f(\gamma(t))x'(t) + i f(\gamma(t))y'(t) \right)}_{f(\gamma(t))\gamma'(t)} dt.$$

In fact, if you also write $d\bar{z} = dx - i dy$, you can write any integral

$$\int_{\gamma} P dx + Q dy \quad \text{as} \quad \int_{\gamma} F dz + G d\bar{z}$$

and vice versa (exercise).